arXiv:gr-qc/9611050v2 3 Dec 1996

SNUTP 96-117

4-Dimensional Kaluza-Klein Approach to General Relativity in the (2,2)-Splitting of Spacetimes

J.H. Yoon*
Department of Physics
Kon-Kuk University, Seoul 143-701, Korea

Abstract

We propose a Kaluza-Klein approach to general relativity of 4-dimensional spacetimes. This approach is based on the (2,2)-splitting of a generic 4-dimensional spacetime, which is viewed as a local product of a (1+1)-dimensional base manifold and a 2-dimensional fibre space. In this Kaluza-Klein formalism we find that general relativity of 4-dimensional spacetimes can be interpreted in a natural way as a (1+1)-dimensional gauge theory. In a gauge where the (1+1)-dimensional metric can be written as the Polyakov metric, the action is describable as a (1+1)-dimensional Yang-Mills type gauge theory action coupled to a (1+1)-dimensional non-linear sigma field and a scalar field, subject to the constraint equations associated with the diffeomorphism invariance of the (1+1)-dimensional base manifold. Diffeomorphisms along the fibre directions show up as the Yang-Mills type gauge symmetries, giving rise to the Gauss-law constraints. We also present the Einstein's equations in the Polyakov gauge and discuss them from the (1+1)-dimensional gauge theory point of view. Finally we show that this Kaluza-Klein formalism is closely related to the null hypersurface formalism of general relativity.

PACS numbers: 04.20.-q, 04.20.Cv, 04.20.Fy

Typeset using REVTEX

---

*Electronic address: yoonjh@cosmic.konkuk.ac.kr

I. INTRODUCTION

The exact correspondence of the Euclidean self-dual Einstein's equations to the equations of motion of 2-dimensional non-linear sigma models with the target space as the area-preserving diffeomorphism of 2-surface [1] has inspired us to look further into the intriguing question whether the full-fledged general relativity of 4-dimensional spacetimes can be also formulated as a certain (1+1)-dimensional field theory. Recently we have shown that such a description is indeed possible, by rewriting the Einstein-Hilbert action of general relativity of 4-dimensional spacetimes as a 4-dimensional Kaluza-Klein (KK) action [2–4] in the framework of the (2,2)-decomposition of spacetimes. In this KK approach, the 4-dimensional spacetime is viewed, at least for a finite range of spacetime, as a locally fibred manifold that consists of a (1+1)-dimensional base manifold and a 2-dimensional "auxiliary" fibre space. Remarkably, it turns out that the KK approach, which is often thought a means of dimensional reduction to 4-dimensional spacetimes from higher dimensions via isometries, can be made a useful tool for studying general relativity of a generic spacetime of 4-dimensions from the perspective of lower dimensional spacetimes such as the (1+1)-dimensional spacetime, for instance.

There are certain advantages of this 4-dimensional KK approach to general relativity in the (2,2)-splitting, which led us to develop this formalism. Here we list a few of them. First of all, in (1+1)-dimensions, there exist a number of field theoretic methods recently developed thanks to the string-related theories. Hopefully the rich mathematical methods suitable for (1+1)-dimensional field theories might also prove useful in studying certain aspects of general relativity of 4-dimensional spacetimes, provided that a sensible (1+1)-dimensional field theory description of such spacetimes is available. In fact, in this KK approach based on the (2,2)-splitting of spacetime, it will be seen that general relativity can be understood in a natural way as a (1+1)-dimensional Yang-Mills type gauge theory coupled to a (1+1)-dimensional non-linear sigma field and a scalar field, with built-in symmetries and interactions prescribed by the diffeomorphism invariance of the associated 4-dimensional spacetimes.

Second, this Yang-Mills type gauge theory description of general relativity simplifies considerably certain issues concerned with the constraints. In Yang-Mills type theories, it is well-known that the Gauss-law constraints associated with the Yang-Mills gauge invariance can be made trivial, if we consider physical, gauge invariant quantities only. Thus, in principle, one might expect that the problem of solving constraints of general relativity could be made *trivial*, at least for some of them, if such a gauge theory description is possible. In the (2,2)-KK approach, the relevant constraints that can be treated this way are those associated with the group of diffeomorphisms of the 2-dimensional fibre space, *not* the entire 4-dimensional spacetime. Of course the remaining two constraints must be studied separately [5].

Third, in the null hypersurface formalism, it is well-known that the exact gravitational wave degrees of freedom reside in the conformal 2-geometry of the wave front 2-surface, transverse to the propagation direction of the gravitational wave. This identification of the exact gravitational degrees of freedom has been particularly useful in the analysis of gravitational waves in the asymptotically flat spacetimes [6,7], and in the asymptotic quantization [8–10] and in the canonical analysis of the null hypersurface formalism [11–13], in the hope of getting quantum theory of gravity by quantizing the true physical degrees of freedom of gravitational field. It turns out, as we shall see later, that the (2,2)-KK formalism is in essence a *novel* reinterpretation of the null hypersurface formalism of 4-dimensional vacuum general relativity from a (1+1)-dimensional point of view, using the Kaluza-Klein ideas. Therefore we expect that this (2,2)-KK formalism will be most appropriate for studying problems involving gravitational waves such as scatterings of gravitational waves, and quantizing gravitational waves, or perhaps for quantum gravity.

Fourth, it is well-known that the null hypersurface formalism, particularly the Newman-Penrose formalism, is outstanding as a method of finding exact solutions of the Einstein's equations. We expect that this feature will also survive in the (2,2)-KK formalism, in view of the manifest correspondence of both formalisms, as is shown in section V and VI.

Last, but not the least, this (2,2)-KK formalism brings about *unusual* conceptual issues related to the interpretation of general relativity. Namely, this (1+1)-dimensional description of 4-dimensional spacetimes allows us to *forget* about the 4-dimensional spacetime picture of general relativity: instead, it enables us to study the subject much the way as we study Yang-Mills type gauge theories coupled to matter fields in (1+1)-dimensions, where the Yang-Mills gauge symmetry is an infinite dimensional group of diffeomorphisms of the 2-dimensional fibre space. Apart from the technical questions such as how to deal with such an infinite dimensional group of diffeomorphisms, this (1+1)-dimensional field theory description certainly puts 4-dimensional spacetime physics into a completely *new* perspective, from which a number of interesting speculations on general relativity of 4-dimensional spacetimes could be put forward.

In view of these advantages of the KK formalism in the (2,2)-splitting and the related null hypersurface formalism, it therefore seems worth exploring the (2,2)-KK formalism carefully in order to see what could be learned more about general relativity of 4-dimensional spacetimes. This paper is organized as follows. In section II, we shall outline the general (2,2)-decomposition of 4-dimensional spacetimes, and introduce the KK variables *without* assuming any spacetime Killing vector fields. It will be seen that the Lie derivative, rather than the metric-compatible covariant derivative, appears naturally in this formalism. We defer the details of computations to Appendix B.

In section III, we shall introduce a coordinate condition, known as the Polyakov gauge in the literatures on (1+1)-dimensional field theories. In the Polyakov gauge, the scalar curvature of 4-dimensional spacetime is found to assume a surprisingly simple form that permits a natural (1+1)-dimensional field theoretic interpretation.

In section IV, we shall present the Einstein's equations in the Polyakov gauge in terms of the KK variables. The natural time coordinate in this formalism is found to be the *affine* parameter of the null geodesics. It is worth noting that one of the Einstein's equations in this formalism can be written in a form of the Schrödinger equation, reminiscent of the Brill wave equation.

In section V, a review of the null hypersurface formalism is given, and the relation between the KK variables and the null tetrads is explicitly shown. Then, in section VI, we shall establish the correspondence between the KK field equations and the Einstein's equations, using the well-known classification of the Einstein's equations in the null hypersurface formalism. We summarize this paper in section VII.

In Appendix A, we shall find the transformation properties of the KK variables *without* assuming any spacetime Killing vector fields, and identify the KK scalar, tensor and gauge fields in the absence of such Killing vector fields. In Appendix B, we shall present the prescription of computing the scalar curvature of 4-dimensional spacetimes, using the KK variables. The intermediate steps of computations are also presented, in the hope that computations can be relatively easily checked. We shall also present the surface terms in the action integral for future references, although we shall not use them in this paper. In Appendix C, it is shown that κ^2, which appears in the discussion of the Einstein's equations in section IV, is positive definite.

II. KALUZA-KLEIN FORMALISM IN THE (2,2)-SPLITTING OF SPACETIMES

In this section we shall describe the general (2,2)-decomposition of 4-dimensional spacetimes from the KK perspective, where spacetime is viewed as a *locally* fibred manifold, i.e. a local direct product of a (1+1)-dimensional *base* manifold M_{1+1} and a 2-dimensional *fibre* space N_2, for which we introduce two pairs of basis vector fields $\partial_\mu = \partial/\partial x^\mu$ $(\mu = 0, 1)$ and $\partial_a = \partial/\partial y^a (a = 2, 3)$, respectively. Let us denote by $\hat{\partial}_\mu$ the vector fields tangent to the horizontal space, orthogonal to the fibre space N_2. These horizontal vector fields can be written as linear combinations of ∂_μ and ∂_a,

$$\hat{\partial}_\mu = \partial_\mu - A_\mu^{\ a} \partial_a, \tag{2.1}$$

where $A_\mu^{\ a}$ are the coefficient functions of the linear combinations. Let us denote by $\gamma_{\mu\nu}$ the metric on the horizontal space, and by ϕ_{ab} the metric on N_2, respectively. Then, without loss of generality, the line element of the 4-dimensional spacetime can be written as [14,15]

$$ds^2 = \gamma_{\mu\nu} dx^\mu dx^\nu + \phi_{ab} \Big(dy^a + A_\mu^{\ a} dx^\mu \Big) \Big(dy^b + A_\nu^{\ b} dx^\nu \Big), \tag{2.2}$$

or in components,

$$g_{AB} = \begin{pmatrix} \gamma_{\mu\nu} + \phi_{ab} A_\mu^{\ a} A_\nu^{\ b} & \phi_{ab} A_\mu^{\ b} \\ \phi_{ab} A_\nu^{\ a} & \phi_{ab} \end{pmatrix}, \tag{2.3}$$

where $A = (\mu, a)$. Formally this metric looks similar to the "dimensionally reduced" metric in standard KK theories, but in fact it is quite different. In the standard KK reduction we normally assume isometries, and the dimensional reduction is made by projection along the directions generated by these isometries. There the fields $A_\mu^{\ a}$ are identified as the KK gauge fields associated with the isometry group. In this paper, we do *not* assume such isometries: nevertheless, it turns out that the KK idea still works. To see this, let us consider the following coordinate transformations of N_2 (diffN_2), while keeping x^μ constant,

$$\begin{aligned} y'^{a} &= y'^{a}(x, y), \\ x'^{\mu} &= x^\mu, \end{aligned} \tag{2.4}$$

where $y'^{a}(x, y)$ is an arbitrary function of x^μ and y^a. Under these transformations, we find the metric coefficients transform as (see Appendix A),

$$\phi'_{ab}(x',y') = \frac{\partial y^c}{\partial y'^a}\frac{\partial y^d}{\partial y'^b}\phi_{cd}(x,y), \tag{2.5}$$

$$A_\mu{}'^a(x',y') = \frac{\partial y'^a}{\partial y^c}A_\mu{}^c(x,y) - \partial_\mu y'^a(x,y), \tag{2.6}$$

$$\gamma'_{\mu\nu}(x',y') = \gamma_{\mu\nu}(x,y), \tag{2.7}$$

which follows from the invariance of the line element (2.2) under the diffN_2 transformations. Under the corresponding infinitesimal transformation

$$\begin{aligned} \delta y^a &= \xi^a(x,y) \qquad (0(\xi^2) \ll 1), \\ \delta x^\mu &= 0, \end{aligned} \tag{2.8}$$

where $\xi^a(x,y)$ is an arbitrary function of x^μ and y^a, the fields transform as

$$\begin{aligned} \delta\phi_{ab}(x,y) &= -\xi^c\partial_c\phi_{ab} - (\partial_a\xi^c)\phi_{cb} - (\partial_b\xi^c)\phi_{ac} \\ &= -\pounds_\xi\phi_{ab} \\ &= -[\xi,\phi]_{\mathrm{L}ab}, \end{aligned} \tag{2.9}$$

$$\begin{aligned} \delta A_\mu{}^a(x,y) &= -\partial_\mu\xi^a + A^c_\mu\partial_c\xi^a - \xi^c\partial_c A_\mu{}^a \\ &= -\partial_\mu\xi^a + \pounds_{A_\mu}\xi^a \\ &= -\partial_\mu\xi^a + [A_\mu,\ \xi]^a_{\mathrm{L}}, \end{aligned} \tag{2.10}$$

$$\begin{aligned} \delta\gamma_{\mu\nu}(x,y) &= -\xi^c\partial_c\gamma_{\mu\nu} \\ &= -\pounds_\xi\gamma_{\mu\nu} \\ &= -[\xi,\gamma_{\mu\nu}]_{\mathrm{L}}. \end{aligned} \tag{2.11}$$

Here $\xi = \xi^a(x,y)\partial_a$ and $A_\mu = A_\mu{}^a(x,y)\partial_a$ are vector fields tangent to N_2, and the brackets with the subscript $_{\mathrm{L}}$ stand for the *Lie* derivatives[1]. Associated with this diffN_2 transformation, the diffN_2-covariant derivative D_μ is defined naturally as

$$D_\mu = \partial_\mu - [A_\mu,\ \ ]_{\mathrm{L}}. \tag{2.12}$$

With this definition, it follows that

$$\delta A_\mu{}^a = -D_\mu\xi^a, \tag{2.13}$$

which clearly indicates that $A_\mu{}^a$ are the gauge fields valued in the infinite dimensional Lie algebra of diffN_2. Moreover, a *covariant* derivative of the field ϕ_{ab} is defined as

$$\begin{aligned} D_\mu\phi_{ab} &= \partial_\mu\phi_{ab} - [A_\mu,\phi]_{\mathrm{L}ab} \\ &= \partial_\mu\phi_{ab} - A_\mu{}^c\partial_c\phi_{ab} - (\partial_a A_\mu{}^c)\phi_{bc} - (\partial_b A_\mu{}^c)\phi_{ac}. \end{aligned} \tag{2.14}$$

The field strength $F_{\mu\nu}{}^a$ corresponding to $A_\mu{}^a$ can now be defined as

---

[1] Here the Lie derivative is defined such that it acts on the fibre space indices $a,b,c,\cdots$ only.

$$\begin{aligned} F_{\mu\nu}{}^{a} &= \partial_\mu A_\nu{}^{a} - \partial_\nu A_\mu{}^{a} - [A_\mu, A_\nu]_{\mathrm{L}}^{a} \\ &= \partial_\mu A_\nu{}^{a} - \partial_\nu A_\mu{}^{a} - A_\mu{}^{c}\partial_c A_\nu{}^{a} + A_\nu{}^{c}\partial_c A_\mu{}^{a}. \end{aligned} \tag{2.15}$$

As is shown in Appendix A, $D_\mu \phi_{ab}$ and $F_{\mu\nu}{}^{a}$ transform *covariantly* under the infinitesimal transformation (2.8),

$$\begin{aligned} \delta(D_\mu\phi_{ab}) &= -\xi^c\partial_c(D_\mu\phi_{ab}) - (\partial_a\xi^c)(D_\mu\phi_{bc}) - (\partial_b\xi^c)(D_\mu\phi_{ac}) \\ &= -[\xi, D_\mu\phi]_{\mathrm{L}ab}, \end{aligned} \tag{2.16}$$

$$\begin{aligned} \delta F_{\mu\nu}{}^{a} &= -\xi^b\partial_b F_{\mu\nu}{}^{a} + F_{\mu\nu}{}^{b}\partial_b\xi^a \\ &= -[\xi, F_{\mu\nu}]_{\mathrm{L}}^{a}. \end{aligned} \tag{2.17}$$

It must be marked here that, in the (2,2)-KK formalism, the Lie derivative, rather than the covariant derivative, appears naturally[2]. Particularly, in the defining equation (2.12) of the diffN_2-covariant derivative, the Lie derivative along the vector fields A_μ is formally identical to the Lie commutator through which the *minimal* couplings to gauge fields are introduced in conventional gauge theories. Together with the fact that the Lie derivative and the Lie commutator, which is usually represented by a finite dimensional matrix commutator, share the same Lie algebraic properties at the abstract level, this observation suggests that the Lie derivative, associated with diffN_2 here, may be viewed as a natural infinite dimensional generalization of the finite dimensional matrix commutator. The appearance of an infinite dimensional symmetry such as diffN_2 is not that surprising, since in general relativity the underlying symmetry is the infinite dimensional group of diffeomorphisms of 4-dimensional spacetime. It is the diffN_2 symmetry, the subgroup of diffeomorphisms of 4-dimensional spacetime, that shows up as a local gauge symmetry[3] of the Yang-Mills type, where the usual minimal couplings are replaced by the Lie derivatives along the vector fields A_μ. We also note that ϕ_{ab} and $\gamma_{\mu\nu}$ transform as a tensor field and scalar fields under the diffN_2 transformations, respectively, as can be seen from (2.9) and (2.11). This implies that the (2,2)-KK formalism can be made a viable method of studying general relativity from the standpoint of (1+1)-dimensional gauge theories with an infinite dimensional gauge symmetry of the Yang-Mills type.

The prescription to obtain the scalar curvature of 4-dimensional spacetime in terms of the KK variables is given in Appendix B. Here we present the result only. It is given by

$$\begin{aligned} R = {}& \gamma^{\mu\nu}\tilde{R}_{\mu\nu} + \phi^{ac}R_{ac} + \frac{1}{4}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b} + \frac{1}{4}\gamma^{\mu\nu}\phi^{ab}\phi^{cd}\Big\{(D_\mu\phi_{ac})(D_\nu\phi_{bd}) \\ & -(D_\mu\phi_{ab})(D_\nu\phi_{cd})\Big\} + \frac{1}{4}\phi^{ab}\gamma^{\mu\nu}\gamma^{\alpha\beta}\Big\{(\partial_a\gamma_{\mu\alpha})(\partial_b\gamma_{\nu\beta}) - (\partial_a\gamma_{\mu\nu})(\partial_b\gamma_{\alpha\beta})\Big\} \\ & +(\hat{\nabla}_\mu + \hat{\Gamma}_{c\mu}{}^{c})j^\mu + (\hat{\nabla}_a + \hat{\Gamma}_{\alpha a}{}^{\alpha})j^a, \end{aligned} \tag{2.18}$$

---

[2] There are occasions when the Lie derivative becomes the natural derivative operator. For instance, at null infinity $\mathcal{I}^+$, the natural derivative operator is the Lie derivative, rather than the covariant derivative ∇_A, because the metric on $\mathcal{I}^+$ is degenerate with signature $(0,+,+)$ [16].

[3] Here local means local in the (1+1)-dimensional base manifold M_{1+1}.

using the notations defined in Appendix B. The Einstein-Hilbert action is given by the integral

$$S = \int d^2x d^2y \sqrt{-\gamma}\sqrt{\phi}\, R, \tag{2.19}$$

where $\gamma = \det \gamma_{\mu\nu}$ and $\phi = \det \phi_{ab}$. One can easily recognize that this action is in a form of a (1+1)-dimensional field theory action. In geometrical terms the above action can be understood as follows. First, $\gamma^{\mu\nu}\tilde{R}_{\mu\nu}$ is the scalar curvature[4] of the horizontal 2-space normal to the fibre space N_2. Second, $\phi^{ac}R_{ac}$ is the scalar curvature of N_2. When integrated over N_2, it becomes a number proportional to the Euler characteristics of the fibre space N_2. Namely, by the Gauss-Bonnet theorem, we have

$$\int d^2y \sqrt{\phi}\, \phi^{ac}R_{ac} = -4\pi\chi, \tag{2.20}$$

where χ is the Euler characteristics[5]. Third, apart from the last two terms containing j^μ and j^a, which are surface terms as is shown in Appendix B, the remaining terms in (2.18) are the *extrinsic* terms, telling how the two 2-surfaces are embedded into the enveloping 4-dimensional spacetime. They are the Yang-Mills term, and a generic non-linear sigma model and a scalar field term. Thus, the Einstein-Hilbert action is describable as a (1+1)-dimensional Yang-Mills type gauge theory coupled to a (1+1)-dimensional non-linear sigma field and scalar fields of generic types, with intrinsic curvature terms of the two 2-surfaces. The associated Yang-Mills gauge symmetry is the group of diffeomorphisms of N_2.

Although the action (2.19) already has several features of gauge theories as described above, it appears rather complicated that its practical value is a bit obscure. In the following section, we will find a simple action integral, more faithful to the spirit of gauge theories, by suitably gauge-fixing some of the spacetime diffeomorphisms. This gauge-fixing brings out the Yang-Mills aspects of general relativity more clearly, as we shall see shortly.

III. KALUZA-KLEIN FORMALISM IN POLYAKOV GAUGE

In the previous section we presented the Einstein-Hilbert action in the general $(2,2)$-splitting of spacetimes in terms of the KK variables. The most noticeable features of the above formalism, among others, are that the diffN_2 symmetry appears as the *built-in* local gauge symmetry of the Yang-Mills type and that $A_\mu^{\ a}$ can be identified as the associated gauge fields. Let us recall that we can always gauge away two functions in the metric $\gamma_{\mu\nu}$ by a suitable coordinate transformation, so that the metric can be written as, at least locally,

---

[4] It may be worth noting that, from the standpoint of the (1+1)-dimensional base manifold M_{1+1}, the scalar curvature of the horizontal space looks like a "gauged" scalar curvature in that it has the diffN_2-valued gauge fields coupled to $\gamma_{\mu\nu}$ in the curvature formula (B34).

[5] Note that the extra minus sign in (2.20) is due to our sign convention of curvature formula, which is the minus of more conventional one [15]. For instance χ is 2 for S^2.

$$\gamma_{\mu\nu} = \begin{pmatrix} -2h & -1 \\ -1 & 0 \end{pmatrix}, \qquad \gamma^{\mu\nu} = \begin{pmatrix} 0 & -1 \\ -1 & 2h \end{pmatrix}. \tag{3.1}$$

Let $(u,\, v)$ be a new coordinate system on M_{1+1}, in which the metric $\gamma_{\mu\nu}$ is of the form (3.1). In the coordinate $(u,\, v,\, y^a)$, the line element (2.2) in section II becomes

$$ds^2 = -2\, du dv - 2h\, du^2 + \phi_{ab}\Big(dy^a + A_+{}^a du + A_-{}^a dv\Big)\Big(dy^b + A_+{}^b du + A_-{}^b dv\Big). \tag{3.2}$$

It is easy to see that the remaining fields ϕ_{ab}, $A_\pm{}^a$, and h still transform as a tensor field, gauge fields, and a scalar field, respectively, under the diffN_2 transformations. Under the diffN_2 transformations

$$y^{'a} = y^{'a}(u,v,y), \qquad u' = u, \qquad v' = v, \tag{3.3}$$

we find that ϕ_{ab}, $A_\pm{}^a$, and h transform as

$$\phi'_{ab}(u',v',y') = \frac{\partial y^c}{\partial y^{'a}}\frac{\partial y^d}{\partial y^{'b}}\phi_{cd}(u,v,y), \tag{3.4}$$

$$A_\pm^{'a}(u',v',y') = \frac{\partial y^{'a}}{\partial y^c}A_\pm{}^c(u,v,y) - \partial_\pm y^{'a}(u,v,y), \tag{3.5}$$

$$h'(u',v',y') = h(u,v,y), \tag{3.6}$$

where $\partial_+ = \partial/\partial u$ and $\partial_- = \partial/\partial v$. Under the corresponding infinitesimal transformations

$$\begin{aligned} &\delta y^a = \xi^a(u,v,y) \qquad (\mathrm{O}(\xi^2) \ll 1), \\ &\delta u = \delta v = 0, \end{aligned} \tag{3.7}$$

they transform as

$$\delta\phi_{ab}(u,v,y) = -[\xi,\phi]_{\mathrm{L}ab}, \tag{3.8}$$

$$\delta A_\pm{}^a(u,v,y) = -D_\pm\xi^a = -\partial_\pm\xi^a + [A_\pm,\xi]_{\mathrm{L}}^a, \tag{3.9}$$

$$\delta h(u,v,y) = -[\xi,h]_{\mathrm{L}}. \tag{3.10}$$

Here the Lie derivatives are defined as before,

$$[\xi,\phi]_{\mathrm{L}ab} = \xi^c\partial_c\phi_{ab} + (\partial_a\xi^c)\phi_{cb} + (\partial_b\xi^c)\phi_{ac}, \tag{3.11}$$

$$[A_\pm,\xi]_{\mathrm{L}}^a = A_\pm{}^c\partial_c\xi^a - \xi^c\partial_c A_\pm{}^a, \tag{3.12}$$

$$[\xi,h]_{\mathrm{L}} = \xi^c\partial_c h. \tag{3.13}$$

This observation tells us two things. First, diffN_2 is (a part of) the residual symmetry of the line element (3.2), and the presence of this residual symmetry allows us to reduce the metric further. To see this, let us consider the following infinitesimal transformation of the gauge field $A_-{}^a$,

$$A_-^{'a} = A_-{}^a - \partial_-\xi^a + [A_-,\xi]_{\mathrm{L}}^a. \tag{3.14}$$

Let ξ^a be a solution of the equation $A_-^{\prime a} = 0$. This equation defines a system of the first-order linear partial differential equations [17] for ξ^a. The existence of a solution of the above equation implies that we can always choose the gauge condition

$$A_-^{\prime a} = 0, \tag{3.15}$$

at least locally. When this condition is satisfied, the line element can be written as

$$ds^2 = -2\,dudv - 2h\,du^2 + \phi_{ab}\Big(dy^a + A_+^{\ a}du\Big)\Big(dy^b + A_+^{\ b}du\Big), \tag{3.16}$$

and its inverse metric is given by

$$g^{AB} = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 2h & A_+^{\ a} \\ 0 & A_+^{\ a} & \phi^{ab} \end{pmatrix}. \tag{3.17}$$

The coordinate system in which the metric is of the form (3.16) shall be referred to as the Newman-Unti coordinate [18], for reasons that will become clear in section V. Here we simply note that the $u =$ constant surface in (3.16) is a degenerate null hypersurface. The null vector field that lies in the null hypersurface is $\partial/\partial v$, since it has a zero norm. In each null hypersurface labeled by u, the $v =$ constant section defines a 2-dimensional spacelike surface N_2 transverse to the vector field $\partial/\partial v$. In order to exhibit the diffN_2 symmetry manifest, however, we shall work mostly with the line element (3.2) without picking up the condition (3.15), although we can always enforce this condition whenever necessary. Second, it is clear that diffN_2 can be still viewed as the local gauge symmetry of the Yang-Mills type, in view of the transformation properties of the fields.

The scalar curvature of 4-dimensional spacetime in the Polyakov gauge (3.1) can be written in a remarkably simple form. In order to see this, let us compute each term in (2.18) in section II using the Polyakov metric. The formulae for the connection coefficients $\hat{\Gamma}_{\mu\nu}^{\ \ \ \alpha}$ are given in (B14) in Appendix B, which become, in the Polyakov gauge,

$$\begin{aligned} &\hat{\Gamma}_{++}^{\ \ \ +} = -\hat{\Gamma}_{+-}^{\ \ \ -} = -\hat{\Gamma}_{-+}^{\ \ \ -} = -D_-h, \\ &\hat{\Gamma}_{++}^{\ \ \ -} = D_+h + 2hD_-h, \end{aligned} \tag{3.18}$$

and zero otherwise. Here $D_\pm h$, the diffN_2-covariant derivatives of the scalar field h, are defined as

$$\begin{aligned} D_\pm h &= \partial_\pm h - [A_\pm, h] \\ &= \partial_\pm h - A_\pm^{\ a}\partial_a h. \end{aligned} \tag{3.19}$$

The Ricci tensor $\tilde{R}_{\mu\nu}$ of the horizontal 2-space is given in (B34) in Appendix B. They become, in the Polyakov gauge,

$$\begin{aligned} &\tilde{R}_{--} = 0, \\ &\tilde{R}_{+-} = \tilde{R}_{-+} = -D_-^2 h, \\ &\tilde{R}_{++} = D_+D_-h - D_-D_+h - 2hD_-^2 h \\ &\qquad = -F_{+-}^{\ \ \ a}\partial_a h - 2hD_-^2 h, \end{aligned} \tag{3.20}$$

where in the last line we used the identity

$$D_+ D_- h - D_- D_+ h = -F_{+-}{}^a \partial_a h, \tag{3.21}$$

where

$$F_{+-}{}^a = \partial_+ A_-{}^a - \partial_- A_+^a - [A_+, A_-]_{\rm L}^a. \tag{3.22}$$

Therefore we have

$$\begin{aligned} \gamma^{\mu\nu} \tilde{R}_{\mu\nu} &= 2\gamma^{+-} \tilde{R}_{+-} \\ &= 2D_-^2 h. \end{aligned} \tag{3.23}$$

Let us further notice that

$$\gamma^{\mu\nu} \partial_a \gamma_{\mu\nu} = \frac{2}{\sqrt{-\gamma}} \partial_a \sqrt{-\gamma} = 0, \tag{3.24}$$

since $\sqrt{-\gamma} = 1$. Also we have

$$\gamma^{\mu\nu} \gamma^{\alpha\beta} (\partial_a \gamma_{\mu\alpha})(\partial_b \gamma_{\nu\beta}) = 0, \tag{3.25}$$

since $\gamma^{++} = 0$ and $\gamma^{+-} = -1$. Therefore it follows that

$$\frac{1}{4} \phi^{ab} \gamma^{\mu\nu} \gamma^{\alpha\beta} \Big\{ (\partial_a \gamma_{\mu\alpha})(\partial_b \gamma_{\nu\beta}) - (\partial_a \gamma_{\mu\nu})(\partial_b \gamma_{\alpha\beta}) \Big\} = 0, \tag{3.26}$$

provided that the Polyakov gauge is chosen. The terms containing $D_\pm \phi_{ab}$ must also be written in this gauge. The diffN_2-covariant derivative of ϕ_{ab} in the Polyakov gauge is defined as

$$\begin{aligned} D_\pm \phi_{ab} &= \partial_\pm \phi_{ab} - [A_\pm, \phi]_{{\rm L}ab} \\ &= \partial_\pm \phi_{ab} - A_\pm^c \partial_c \phi_{ab} - (\partial_a A_\pm^c) \phi_{cb} - (\partial_b A_\pm^c) \phi_{ac}. \end{aligned} \tag{3.27}$$

It is well-known that, in the null hypersurface formalism of general relativity, the transverse 2-metric with a unit determinant, i.e. the conformal 2-metric [7,19–21] represents the two physical degrees of freedom of gravitational field[6]. Let us therefore decompose the metric ϕ_{ab} of the fibre space N_2 into a volume field ϕ and a conformal 2-metric ρ_{ab},

$$\phi_{ab} = \sqrt{\phi}\ \rho_{ab} = {\rm e}^\sigma \rho_{ab}, \tag{3.28}$$

where ρ_{ab} satisfies the condition

$$\det \rho_{ab} = 1, \tag{3.29}$$

---

[6] The gauge in which the metric is of the form (3.2) may be viewed as an analog of the Coulomb gauge in Maxwell's theory, where the physical degrees of freedom are the transverse traceless vector potentials, since a finite generalization of traceless condition is a unit determinant condition.

so that $\phi = \det \phi_{ab}$. The field σ is related to ϕ by the equation

$$\sigma = \frac{1}{2} \ln \phi. \tag{3.30}$$

Notice that ρ_{ab} is a tensor density with the weight $+1$, and that ϕ is a scalar density with the weight -2 with respect to the diffN_2 transformation, respectively. Therefore the diffN_2-covariant derivatives[7] of ρ_{ab} and ϕ are naturally given by

$$\begin{aligned} D_\pm \rho_{ab} &= \partial_\pm \rho_{ab} - [A_\pm, \rho]_{\mathrm{L}ab} \\ &= \partial_\pm \rho_{ab} - A_\pm{}^c \partial_c \rho_{ab} - (\partial_a A_\pm{}^c)\rho_{cb} - (\partial_b A_\pm{}^c)\rho_{ac} + (\partial_c A_\pm{}^c)\rho_{ab}, \end{aligned} \tag{3.31}$$

$$\begin{aligned} D_\pm \phi &= \partial_\pm \phi - [A_\pm, \phi]_{\mathrm{L}} \\ &= \partial_\pm \phi - A_\pm{}^a \partial_a \phi - 2(\partial_a A_\pm{}^a)\phi, \end{aligned} \tag{3.32}$$

respectively. The diffN_2-covariant of σ follows automatically from (3.32),

$$\begin{aligned} D_\pm \sigma &= \partial_\pm \sigma - [A_\pm, \sigma]_{\mathrm{L}} \\ &= \partial_\pm \sigma - A_\pm{}^a \partial_a \sigma - \partial_a A_\pm{}^a. \end{aligned} \tag{3.33}$$

Using the identity

$$\rho^{ab} D_\pm \rho_{ab} = 0, \tag{3.34}$$

which is as a trivial consequence of (3.29) and (3.31), we find that

$$\begin{aligned} \phi^{ab} D_\pm \phi_{ab} &= \mathrm{e}^{-\sigma} \rho^{ab} \Big\{ \mathrm{e}^{\sigma} (D_\pm \sigma)\rho_{ab} + \mathrm{e}^{\sigma} D_\pm \rho_{ab} \Big\} \\ &= 2 D_\pm \sigma. \end{aligned} \tag{3.35}$$

Let us notice that we also have

$$\begin{aligned} &\phi^{ab}\phi^{cd}(D_\pm \phi_{ac})(D_- \phi_{bd}) \\ &= \mathrm{e}^{-2\sigma} \rho^{ab}\rho^{cd} \Big\{ \mathrm{e}^{\sigma}(D_\pm \sigma)\rho_{ac} + \mathrm{e}^{\sigma}(D_\pm \rho_{ac}) \Big\} \Big\{ \mathrm{e}^{\sigma}(D_- \sigma)\rho_{bd} + \mathrm{e}^{\sigma}(D_- \rho_{bd}) \Big\} \\ &= 2(D_\pm \sigma)(D_- \sigma) + \rho^{ab}(D_\pm \sigma)(D_- \rho_{ab}) + \rho^{ab}(D_- \sigma)(D_\pm \rho_{ab}) + \rho^{ab}\rho^{cd}(D_\pm \rho_{ac})(D_- \rho_{bd}) \\ &= 2(D_\pm \sigma)(D_- \sigma) + \rho^{ab}\rho^{cd}(D_\pm \rho_{ac})(D_- \rho_{bd}), \end{aligned} \tag{3.36}$$

using the equation (3.34). Therefore the term containing $D_\pm \phi_{ab}$ becomes

$$\begin{aligned} &\frac{1}{4}\gamma^{\mu\nu}\phi^{ab}\phi^{cd} \Big\{ (D_\mu \phi_{ac})(D_\nu \phi_{bd}) - (D_\mu \phi_{ab})(D_\nu \phi_{cd}) \Big\} \\ &= (D_+ \sigma)(D_- \sigma) - \frac{1}{2}\rho^{ab}\rho^{cd}(D_+ \rho_{ac})(D_- \rho_{bd}) - h\Big\{ (D_- \sigma)^2 - \frac{1}{2}\rho^{ab}\rho^{cd}(D_- \rho_{ac})(D_- \rho_{bd}) \Big\}. \end{aligned} \tag{3.37}$$

---

[7] In general, for the tensor density $T_{ab\cdots}$ with the weight w with respect to the diffN_2 transformations, we have $D_\pm T_{ab\cdots} = \partial_\pm T_{ab\cdots} - [A_\pm, T]_{\mathrm{L}ab\cdots}$, which includes the term $+w(\partial_c A_\pm{}^c) T_{ab\cdots}$.

Also the Yang-Mills type term becomes,

$$\frac{1}{4}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b} = -\frac{1}{2}\mathrm{e}^{\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b}. \tag{3.38}$$

The remaining terms are the surface terms when integrated over the spacetime coordinates, as is shown in Appendix B. For completeness, we shall evaluate them in the Polyakov gauge, using the defining equations of $j^{\pm}$ and j^{a},

$$j^{\mu} = \gamma^{\mu\nu}\phi^{ab}D_{\nu}\phi_{ab}, \tag{3.39}$$

$$j^{a} = \phi^{ab}\gamma^{\mu\nu}\partial_{b}\gamma_{\mu\nu}, \tag{3.40}$$

as is given in (B41) and (B42) in Appendix B. From the equation (3.24) we have

$$j^{a} = 0, \tag{3.41}$$

and $j^{\pm}$ are found to be

$$j^{+} = -2D_{-}\sigma, \tag{3.42}$$

$$j^{-} = -2D_{+}\sigma + 4hD_{-}\sigma. \tag{3.43}$$

Therefore the surface term containing $j^{\pm}$ becomes,

$$\begin{aligned}\Big(\hat{\nabla}_{\mu} + \hat{\Gamma}_{c\mu}{}^{c}\Big)j^{\mu} &= \hat{\partial}_{\mu}j^{\mu} + \hat{\Gamma}_{\mu\nu}{}^{\mu}j^{\nu} + \hat{\Gamma}_{c\mu}{}^{c}j^{\mu}\\ &= \hat{\partial}_{+}j^{+} + \hat{\partial}_{-}j^{-} + \hat{\Gamma}_{++}{}^{+}j^{+} + \hat{\Gamma}_{-+}{}^{-}j^{+} + \hat{\Gamma}_{c+}{}^{c}j^{+} + \hat{\Gamma}_{c-}{}^{c}j^{-}\\ &= \Big(\hat{\partial}_{+} + \hat{\Gamma}_{c+}{}^{c}\Big)j^{+} + \Big(\hat{\partial}_{-} + \hat{\Gamma}_{c-}{}^{c}\Big)j^{-},\end{aligned} \tag{3.44}$$

where we used the identities

$$\begin{aligned}&\hat{\Gamma}_{+-}{}^{+} = \hat{\Gamma}_{--}{}^{-} = 0,\\ &\hat{\Gamma}_{++}{}^{+} = -\hat{\Gamma}_{-+}{}^{-} = -D_{-}h.\end{aligned} \tag{3.45}$$

Using the equation

$$\hat{\Gamma}_{c\pm}{}^{c} = D_{\pm}\sigma, \tag{3.46}$$

which follows from (B27) in Appendix B and (3.35), the surface term finally becomes,

$$\begin{aligned}\Big(\hat{\nabla}_{\mu} + \hat{\Gamma}_{c\mu}{}^{c}\Big)j^{\mu} = &-2\Big(\partial_{+} - A_{+}{}^{a}\partial_{a} + D_{+}\sigma\Big)(D_{-}\sigma)\\ &-2\Big(\partial_{-} - A_{-}{}^{a}\partial_{a} + D_{-}\sigma\Big)\Big(D_{+}\sigma - 2hD_{-}\sigma\Big).\end{aligned} \tag{3.47}$$

Thus, putting the equations (3.23), (3.26), (3.37), (3.38), and (3.47) all together into (2.18) in section II, we obtain the scalar curvature[8] $R_{\rm p}$ of 4-dimensional spacetime in the Polyakov gauge. It is given by

---

[8] Here the subscript $_{\rm p}$ means that the quantity was evaluated in the Polyakov gauge.

$$\begin{aligned} R_{\rm p} = R_2 &+ 2D_-^2 h - \frac{1}{2}{\rm e}^{\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} + (D_+\sigma)(D_-\sigma) - \frac{1}{2}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_-\rho_{bd}) \\ &-h\Big\{(D_-\sigma)^2 - \frac{1}{2}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\} - 2\Big(\partial_+ - A_+{}^{a}\partial_a + D_+\sigma\Big)(D_-\sigma) \\ &-2\Big(\partial_- - A_-{}^{a}\partial_a + D_-\sigma\Big)\Big(D_+\sigma - 2hD_-\sigma\Big), \end{aligned} \tag{3.48}$$

where $R_2 = \phi^{ac}R_{ac}$ is the scalar curvature of the fibre space N_2. The integral of $R_{\rm p}$ over the 4-dimensional spacetime, which we denote by $S_{\rm p}$, is given by

$$S_{\rm p} = \int du\, dv\, d^2y\, {\rm e}^{\sigma} R_{\rm p}, \tag{3.49}$$

since $\sqrt{-\gamma} = 1$. Ignoring the surface integrals coming from the last two terms in $R_{\rm p}$, the integral $S_{\rm p}$ can be written as

$$\begin{aligned} S_{\rm p} = \int du\, dv\, d^2y\, \Big[&-\frac{1}{2}{\rm e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} + {\rm e}^{\sigma}(D_+\sigma)(D_-\sigma) - \frac{1}{2}{\rm e}^{\sigma}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_-\rho_{bd}) \\ &+{\rm e}^{\sigma}R_2 + 2{\rm e}^{\sigma}D_-^2 h - h{\rm e}^{\sigma}\Big\{(D_-\sigma)^2 - \frac{1}{2}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\}\Big] \\ &+ \text{surface terms}. \end{aligned} \tag{3.50}$$

Let us now notice that the term ${\rm e}^{\sigma}D_-^2 h$ can be written as

$$\begin{aligned} {\rm e}^{\sigma}D_-^2 h &= -{\rm e}^{\sigma}(D_-\sigma)(D_-h) + D_-\Big({\rm e}^{\sigma}D_-h\Big) \\ &= -{\rm e}^{\sigma}(D_-\sigma)(D_-h) + \partial_-\Big({\rm e}^{\sigma}D_-h\Big) - \partial_a\Big(A_-{}^{a}{\rm e}^{\sigma}D_-h\Big). \end{aligned} \tag{3.51}$$

But the first term in (3.51) becomes

$$\begin{aligned} &-{\rm e}^{\sigma}(D_-\sigma)(D_-h) \\ &= hD_-({\rm e}^{\sigma}D_-\sigma\Big) - D_-\Big(h{\rm e}^{\sigma}D_-\sigma\Big) \\ &= h{\rm e}^{\sigma}(D_-^2\sigma) + h{\rm e}^{\sigma}(D_-\sigma)^2 - \partial_-\Big(h{\rm e}^{\sigma}D_-\sigma\Big) + \partial_a\Big(hA_-{}^{a}{\rm e}^{\sigma}D_-\sigma\Big). \end{aligned} \tag{3.52}$$

Therefore we find that ${\rm e}^{\sigma}D_-^2 h$ becomes

$$\begin{aligned} {\rm e}^{\sigma}D_-^2 h = h{\rm e}^{\sigma}(D_-^2\sigma) &+ h{\rm e}^{\sigma}(D_-\sigma)^2 + \partial_-\Big({\rm e}^{\sigma}D_-h\Big) - \partial_-\Big(h{\rm e}^{\sigma}D_-\sigma\Big) \\ &-\partial_a\Big(A_-{}^{a}{\rm e}^{\sigma}D_-h\Big) + \partial_a\Big(hA_-{}^{a}{\rm e}^{\sigma}D_-\sigma\Big). \end{aligned} \tag{3.53}$$

Thus the integral $S_{\rm p}$ becomes

$$\begin{aligned} S_{\rm p} = \int du\, dv\, d^2y\, \Big[&-\frac{1}{2}{\rm e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} + {\rm e}^{\sigma}(D_+\sigma)(D_-\sigma) - \frac{1}{2}{\rm e}^{\sigma}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_-\rho_{bd}) \\ &+{\rm e}^{\sigma}R_2 + 2h{\rm e}^{\sigma}\Big\{D_-^2\sigma + \frac{1}{2}(D_-\sigma)^2 + \frac{1}{4}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\}\Big] \\ &+\text{surface terms} \\ \equiv \int du\, dv\, d^2y\, &\mathcal{L}_{\rm p} \; + \text{surface terms}, \end{aligned} \tag{3.54}$$

where the integrand $\mathcal{L}_{\rm p}$ is defined as

$$\begin{aligned}\mathcal{L}_{\rm p} = & -\frac{1}{2}{\rm e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} + {\rm e}^{\sigma}(D_{+}\sigma)(D_{-}\sigma) - \frac{1}{2}{\rm e}^{\sigma}\rho^{ab}\rho^{cd}(D_{+}\rho_{ac})(D_{-}\rho_{bd}) \\ & +{\rm e}^{\sigma}R_{2} + 2h{\rm e}^{\sigma}\Big\{D_{-}^{2}\sigma + \frac{1}{2}(D_{-}\sigma)^{2} + \frac{1}{4}\rho^{ab}\rho^{cd}(D_{-}\rho_{ac})(D_{-}\rho_{bd})\Big\}. \qquad (3.55)\end{aligned}$$

This Lagrangian density $\mathcal{L}_{\rm p}$ can be naturally interpreted as describing the Yang-Mills type gauge theory defined on the (1+1)-dimensional base manifold M_{1+1}, interacting with the (1+1)-dimensional "matter" fields ρ_{ab} and σ. The associated local gauge symmetry is the *built-in* diffN_2 symmetry, the group of diffeomorphisms of the fibre space N_2. It must be mentioned here that each term in (3.55) is manifestly diffN_2-invariant, and that the y^a-dependence of each term is completely *hidden* in the Lie derivative. In this sense we may view the fibre space N_2 as a kind of *internal* space[9] as in Yang-Mills theory.

In order to derive the Einstein's equations as the Euler-Lagrange equations of motion in this KK formalism, the Lagrangian density $\mathcal{L}_{\rm p}$ must be implemented with two equations associated with gauge-fixing of the metric $\gamma_{\mu\nu}$ of the horizontal 2-space to the Polyakov metric. Notice that the eight field equations are obtainable from the Lagrangian density $\mathcal{L}_{\rm p}$. In order to find the remaining two equations, we have to vary the general Einstein-Hilbert action (2.19) in section II with respect to the general functions γ_{+-} and γ_{--}, and then use the Polyakov gauge condition $\gamma_{+-} = -1$ and $\gamma_{--} = 0$ at the end of the variations. In the next section we shall present the ten Einstein's equations without giving the details of derivations.

IV. THE EINSTEIN'S EQUATIONS

In this section we present the ten Einstein's equations. The eight equations can be obtained by varying $\mathcal{L}_{\rm p}$ with respect to h, $A_{\pm}{}^{a}$, σ, and ρ_{ab} subject to a unit determinant condition. In order to enforce this condition, we must add to $\mathcal{L}_{\rm p}$ a Lagrange multiplier term

$$\lambda\Big(\det\rho_{ab} - 1\Big), \qquad (4.1)$$

where λ is a Lagrange multiplier to be determined. The eight Einstein's equations are as follows:

$$(a) \quad 2{\rm e}^{\sigma}(D_{-}^{2}\sigma) + {\rm e}^{\sigma}(D_{-}\sigma)^{2} + \frac{1}{2}{\rm e}^{\sigma}\rho^{ab}\rho^{cd}(D_{-}\rho_{ac})(D_{-}\rho_{bd}) = 0, \qquad (4.2)$$

$$\begin{aligned}(b) \quad & D_{-}\Big({\rm e}^{2\sigma}\rho_{ab}F_{+-}{}^{b}\Big) - {\rm e}^{\sigma}\partial_{a}(D_{-}\sigma) - \frac{1}{2}{\rm e}^{\sigma}\rho^{bc}\rho^{de}(D_{-}\rho_{bd})(\partial_{a}\rho_{ce}) + \partial_{b}\Big({\rm e}^{\sigma}\rho^{bc}D_{-}\rho_{ac}\Big) \\ & = 0, \qquad (4.3)\end{aligned}$$

---

[9] The integration over the y^a-coordinates in (3.54) might be viewed as an infinite dimensional generalization of taking finite dimensional *trace* over the fibre space indices in Yang-Mills theories with some finite dimensional gauge symmetries.

$$\begin{aligned}
(c)\quad & -D_+\Big(\mathrm{e}^{2\sigma}\rho_{ab}F_{+-}{}^{b}\Big) - \mathrm{e}^{\sigma}\partial_a(D_+\sigma) - \frac{1}{2}\mathrm{e}^{\sigma}\rho^{bc}\rho^{de}(D_+\rho_{bd})(\partial_a\rho_{ce}) \\
& +\partial_b\Big(\mathrm{e}^{\sigma}\rho^{bc}D_+\rho_{ac}\Big) + 2h\mathrm{e}^{\sigma}\partial_a(D_-\sigma) + h\mathrm{e}^{\sigma}\rho^{bc}\rho^{de}(D_-\rho_{bd})(\partial_a\rho_{ce}) + 2\mathrm{e}^{\sigma}\partial_a(D_-h) \\
& -2\partial_b\Big(h\mathrm{e}^{\sigma}\rho^{bc}D_-\rho_{ac}\Big) = 0, \qquad (4.4) \\
(d)\quad & -2\mathrm{e}^{\sigma}D_-^2h - 2\mathrm{e}^{\sigma}(D_-h)(D_-\sigma) + \mathrm{e}^{\sigma}D_+D_-\sigma + \mathrm{e}^{\sigma}D_-D_+\sigma + \mathrm{e}^{\sigma}(D_+\sigma)(D_-\sigma) \\
& +\frac{1}{2}\mathrm{e}^{\sigma}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_-\rho_{bd}) + \mathrm{e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} - 2h\mathrm{e}^{\sigma}\Big\{D_-^2\sigma + \frac{1}{2}(D_-\sigma)^2 \\
& +\frac{1}{4}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\} = 0, \qquad (4.5) \\
(e)\quad & h\Big\{\mathrm{e}^{\sigma}D_-^2\rho_{ab} - \mathrm{e}^{\sigma}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd}) + \mathrm{e}^{\sigma}(D_-\rho_{ab})(D_-\sigma)\Big\} \\
& -\frac{1}{2}\mathrm{e}^{\sigma}\Big(D_+D_-\rho_{ab} + D_-D_+\rho_{ab}\Big) + \frac{1}{2}\mathrm{e}^{\sigma}\rho^{cd}\Big\{(D_-\rho_{ac})(D_+\rho_{bd}) + (D_-\rho_{bc})(D_+\rho_{ad})\Big\} \\
& -\frac{1}{2}\mathrm{e}^{\sigma}\Big\{(D_-\rho_{ab})(D_+\sigma) + (D_+\rho_{ab})(D_-\sigma)\Big\} \\
& +\mathrm{e}^{\sigma}(D_-\rho_{ab})(D_-h) + \frac{1}{2}\mathrm{e}^{2\sigma}\rho_{ac}\rho_{bd}F_{+-}{}^{c}F_{+-}{}^{d} - \frac{1}{4}\mathrm{e}^{2\sigma}\rho_{ab}\rho_{cd}F_{+-}{}^{c}F_{+-}{}^{d} = 0. \qquad (4.6)
\end{aligned}$$

The remaining two equations associated with the variations with respect to γ_{+-} and γ_{--} are found as follows:

$$\begin{aligned}
(f)\quad & \mathrm{e}^{\sigma}D_+D_-\sigma + \mathrm{e}^{\sigma}D_-D_+\sigma + 2\mathrm{e}^{\sigma}(D_+\sigma)(D_-\sigma) - 2\mathrm{e}^{\sigma}(D_-h)(D_-\sigma) \\
& -\frac{1}{2}\mathrm{e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} - \mathrm{e}^{\sigma}R_2 - h\mathrm{e}^{\sigma}\Big\{(D_-\sigma)^2 - \frac{1}{2}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\} = 0, \qquad (4.7) \\
(g)\quad & -\mathrm{e}^{\sigma}D_+^2\sigma - \frac{1}{2}\mathrm{e}^{\sigma}(D_+\sigma)^2 - \mathrm{e}^{\sigma}(D_-h)(D_+\sigma) + \mathrm{e}^{\sigma}(D_+h)(D_-\sigma) \\
& +2h\mathrm{e}^{\sigma}(D_-h)(D_-\sigma) + \mathrm{e}^{\sigma}F_{+-}{}^{a}\partial_a h - \frac{1}{4}\mathrm{e}^{\sigma}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_+\rho_{bd}) + \partial_a\Big(\rho^{ab}\partial_b h\Big) \\
& +h\Big\{-\mathrm{e}^{\sigma}(D_+\sigma)(D_-\sigma) + \frac{1}{2}\mathrm{e}^{\sigma}\rho^{ab}\rho^{cd}(D_+\rho_{ac})(D_-\rho_{bd}) + \frac{1}{2}\mathrm{e}^{2\sigma}\rho_{ab}F_{+-}{}^{a}F_{+-}{}^{b} + \mathrm{e}^{\sigma}R_2\Big\} \\
& +h^2\mathrm{e}^{\sigma}\Big\{(D_-\sigma)^2 - \frac{1}{2}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd})\Big\} = 0. \qquad (4.8)
\end{aligned}$$

The equations (4.2), (4.3), (4.4), (4.5), and (4.6) are the Euler-Lagrange equations of motion corresponding to the variations in h, $A_+^{\ a}$, $A_-^{\ a}$, σ, and ρ_{ab}, respectively, and the equations (4.7) and (4.8) are the equations of motion of γ_{+-} and γ_{--}, respectively. An inspection of these equations reveals that the four equations (4.4), (4.7), and (4.8) do *not* contain second-order derivative terms in v. This suggests that the affine parameter v may be identified as the *natural* time coordinate in this formalism, and that these four equations are the constraint equations restricting the initial value data on the v = constant hypersurface. Since these four equations are written in the Polyakov gauge, they are expected to generate the residual symmetry[10] [5] of the line element (3.2), i.e. the symmetry that survives after the Polyakov gauge is chosen. Thus, the equations (4.4) generate the diffN_2 symmetry, and the equations (4.7) and (4.8) generate the residual symmetry associated with the reparametrization invariance of the (1+1)-dimensional base manifold M_{1+1}.

---

[10] For the residual symmetry of this kind, see the discussions in [22] for instance.

The remaining equations (4.2), (4.3), (4.5) and (4.6) are the evolution equations since they contain the second-order v-time derivatives. Notice that the equation (4.2) can be written in a form of the Schrödinger equation. Using the identity

$$2\ D_-^2 \mathrm{e}^{\sigma/2} = \mathrm{e}^{\sigma/2}\Big\{D_-^2\sigma + \frac{1}{2}(D_-\sigma)^2\Big\}, \tag{4.9}$$

the equation (4.2) becomes

$$D_-^2\psi + \kappa^2\psi = 0, \tag{4.10}$$

where

$$\begin{aligned} &\psi = \mathrm{e}^{\sigma/2}, \\ &\kappa^2 = \frac{1}{8}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd}) \geq 0. \end{aligned} \tag{4.11}$$

In Appendix C it is shown that κ^2 is positive definite. It is a bilinear combination of the *gravitational current* $j^a{}_b$,

$$j^a{}_b = \rho^{ac}D_-\rho_{cb}, \tag{4.12}$$

formed by the conformal 2-metric ρ_{ab} and its first v-time derivative. Since the true gravitational degrees of freedom are known to reside in the conformal 2-metric ρ_{ab}, we expect that it is the gravitational current $j^a{}_b$ that transports the true gravitational degrees of freedom. More often than not, important physical quantities appear in bilinear combinations of more fundamental quantities. Thus, it is of interest to find what physical quantity κ^2 represents, being a bilinear in the "fundamental" gravitational current[11].

The equation (4.10) is an analog of the Brill wave equation[12] [23], as it is of the type of the Schrödinger equation for a wave function corresponding to a state of zero energy in the potential $-\kappa^2$. Thus σ is a function that can be determined by the potential $-\kappa^2$ up to some integral "constant" function. The generic behavior of solutions of the equation is therefore expected either of the scattering type, or of the bound-state or resonance type, corresponding to the asymptotically flat spacetimes or spatially closed universes, respectively, on a par with the Brill wave equation.

---

[11] In the case of the Brill wave equation, this quantity represents a kind of gravitational energies associated with gravitational waves.

[12] Although the equation (4.10) is a three dimensional P.D.E., it can be made a one dimensional O.D.E. if the gauge condition $A_-{}^a = 0$ is chosen. Unlike the Brill wave equation, the wave function $\mathrm{e}^{\sigma/2}$ here is the square root of the conformal factor of the two dimensional wavefront N_2, the spatial projection of the null hypersurface $u =$ constant.

V. THE NULL HYPERSURFACE FORMALISM

In this section we shall find that the KK formalism presented in previous sections is in fact equivalent to the null hypersurface formalism. This becomes quite obvious if we recognize that the coordinate system in which the Polyakov gauge condition and the extra gauge condition $A_-{}^a = 0$ are simultaneously valid is precisely the Newman-Unti coordinate [18], which is usually adopted in the null hypersurface formalism. We shall begin our discussions by reviewing how the Newman-Unti coordinate is constructed in the null hypersurface formalism. Let us introduce a null vector field l_A, which we may choose to be the minus of the gradient of some scalar function u,

$$l_A = -\nabla_A u. \tag{5.1}$$

From this we can define another scalar function v by the equation

$$l^A \frac{\partial v}{\partial x^A} = 1. \tag{5.2}$$

Solving this equation for l^A, we find that

$$l^A = \frac{\partial x^A}{\partial v}, \tag{5.3}$$

which tells us that v is the (affine) parameter of the null curve generated by l^A. Let us choose $x^+ = u$, $x^- = v$, and $x^a = y^a (a = 2, 3)$ as our coordinates. In this coordinate system, the components of l_A and l^A are given by

$$l_A = -\delta_A{}^+, \qquad l^A = \delta_-{}^A, \tag{5.4}$$

respectively. From the following identity

$$l^A = g^{AB} l_B = \delta_-{}^A, \tag{5.5}$$

we find that g^{A+} is given by

$$g^{A+} = -\delta_-{}^A = (0,\ -1,\ 0,\ 0). \tag{5.6}$$

The coordinate system (u, v, y^a) in which the metric is of the form (5.6) is called the Newman-Unti coordinate, and the metric (3.17) in section III is precisely of this form.

The complete set of the null tetrads $(l,\ n,\ m,\ \bar{m})$ is defined by the following relations

$$\begin{aligned} &l^2 = n^2 = m^2 = \bar{m}^2 = l \cdot m = n \cdot m = 0, \\ &l \cdot n = -1, \qquad m \cdot \bar{m} = 1, \end{aligned} \tag{5.7}$$

where $\bar{m}$ is the complex conjugate of m. In terms of these null tetrads, the metric is given by

$$g^{AB} = -2n^{(A} l^{B)} + 2m^{(A} \bar{m}^{B)}. \tag{5.8}$$

FIGURES


$\frac{\partial}{\partial u}$
u = constant hypersurface
$\frac{\partial}{\partial v}$
$\frac{\partial}{\partial y^a}$
$(a = 2,3)$
(a)
$\frac{\partial}{\partial y^a}(a = 2,3)$
2-dimensional fibre space N_2
$\frac{\partial}{\partial v}$
$\frac{\partial}{\partial u}$
$(1+1)$-dimensional base manifold M_{1+1}
(b)
=

FIG. 1. The relation between the kinematics of the null hypersurface formalism and that of the Kaluza-Klein formalism in the (2,2)-splitting of spacetime is shown.
(a) The null hypersurface formalism: a null hypersurface labelled by u = constant is generated by a null vector field $\partial/\partial v$ and two transverse vector fields $\partial/\partial y^a$.
(b) The Kaluza-Klein formalism: this shows the KK picture of the same geometry from the fibre bundle point of view in the (2,2)-splitting of spacetime. Here the (1+1)-dimensional base manifold M_{1+1} is generated by the vector fields $\partial/\partial u$ and $\partial/\partial v$, and the 2-dimensional fibre space N_2 is generated by the vector fields $\partial/\partial y^a$.

In the Newman-Unti coordinate, these null tetrads can be explicitly constructed as follows. The first null tetrad l is given by

$$l = l^A \frac{\partial}{\partial x^A} = \frac{\partial}{\partial v}, \tag{5.9}$$

which follows from the equation (5.4). The second null tetrad n is chosen such that the following condition

$$l^A n_A = -1 \tag{5.10}$$

holds. The most general form of n compatible with this condition can be written as

$$n = n^A \frac{\partial}{\partial x^A} = \frac{\partial}{\partial u} + U \frac{\partial}{\partial v} + X^a \frac{\partial}{\partial y^a}, \tag{5.11}$$

as can be easily confirmed by inspection. The complex null tetrads m and $\bar{m}$, which satisfy the defining relation (5.7), with l and n given by (5.9) and (5.11), respectively, become

$$m = m^A \frac{\partial}{\partial x^A} = \omega \frac{\partial}{\partial v} + \xi^a \frac{\partial}{\partial y^a}, \tag{5.12}$$

$$\bar{m} = \bar{m}^A \frac{\partial}{\partial x^A} = \bar{\omega} \frac{\partial}{\partial v} + \bar{\xi}^a \frac{\partial}{\partial y^a}, \tag{5.13}$$

where ω and ξ^a are complex quantities[13]. Putting together (5.9), (5.11), (5.12) and (5.13) into (5.8), we find that

$$\begin{aligned} & g^{++} = g^{+a} = 0, \qquad g^{+-} = -1, \\ & g^{--} = -2U + 2\omega\bar{\omega}, \ g^{-a} = -X^a + \omega\bar{\xi}^a + \bar{\omega}\xi^a, \\ & g^{ab} = 2\xi^{(a}\bar{\xi}^{b)}. \end{aligned} \tag{5.14}$$

If we compare these metric coefficients with the ones given in (3.17) in section III, we find that the KK variables can be written in terms of the null tetrads as

$$2h = -2U + 2\omega\bar{\omega}, \tag{5.15}$$

$$A_+^{\ a} = -X^a + \omega\bar{\xi}^a + \bar{\omega}\xi^a, \tag{5.16}$$

$$\phi^{ab} = 2\xi^{(a}\bar{\xi}^{b)}. \tag{5.17}$$

Equivalently, the null tetrads can be written as

$$l = \frac{\partial}{\partial v}, \tag{5.18}$$

$$n = \frac{\partial}{\partial u} + \Big(- h + \omega\bar{\omega} \Big) \frac{\partial}{\partial v} + \Big(- A_+^{\ a} + \omega\bar{\xi}^a + \bar{\omega}\xi^a \Big) \frac{\partial}{\partial y^a}, \tag{5.19}$$

$$m = \omega \frac{\partial}{\partial v} + \xi^a \frac{\partial}{\partial y^a}, \ \text{ and c.c..} \tag{5.20}$$

VI. CLASSIFICATION OF THE FIELD EQUATIONS

It will be instructive to find how the KK field equations (4.2), $\cdots$, (4.8) in section IV are related to the Einstein's equations in the standard null hypersurface formalism. Let us recall that, in the standard null hypersurface formalism, the Einstein's equations are classified into six main equations, one trivial equation, and three supplementary equations. Let G^{AB} be the Einstein tensor. Then the Einstein's equations are classified as

$$(A) \quad \text{main equations} : G^{AB} l_B = 0, \quad G^{AB} m_A m_B = 0 \tag{6.1}$$

$$(B) \quad \text{trivial equation} : G^{AB} m_A \bar{m}_B = 0 \tag{6.2}$$

$$(C) \quad \text{supplementary equations} : G^{AB} n_A m_B = 0, \quad G^{AB} n_A n_B = 0. \tag{6.3}$$

---

[13] Here we may choose the orientation of the complex null tetrad m such that $\omega = 0$, since $\partial / \partial v$ has a zero norm.

Moreover, the Bianchi identity ${G^{AB}}_{;B} = 0$ can be used to prove the following lemma [7]: Suppose that the main equations hold everywhere; then (i) the trivial equation holds everywhere, and (ii) the supplementary equations hold everywhere if they hold at some v = constant hypersurface.

In order to see what these classified equations are like in the Newman-Unti coordinate, we have to find the covariant null tetrads $(l_A,\ n_A,\ m_A,\ \bar{m}_A)$ in the Newman-Unti coordinate first, and then form the above combinations using them. Notice that l_A is already given by the equation (5.4) in section V. To find m_A, we notice that

$$\begin{aligned} m_A &= m^B g_{AB} \\ &= \omega g_{A-} + \xi^a g_{Aa}, \end{aligned} \tag{6.4}$$

since $m^+ = 0$, $m^- = \omega$, and $m^a = \xi^a$. Thus, using g_{AB} in (3.16) in section III, we find that the coefficients of the complex null tetrad m are given by

$$m_+ = -\omega + \xi_a {A_+}^a, \tag{6.5}$$

$$m_- = 0, \tag{6.6}$$

$$m_a = \xi_a, \tag{6.7}$$

where $\xi_a = \phi_{ab}\xi^b$. Similarly, the equation

$$n_A = n^B g_{AB} \tag{6.8}$$

gives

$$n_+ = -2h - U + \Big(\omega\bar{\xi}_a + \bar{\omega}\xi_a\Big){A_+}^a, \tag{6.9}$$

$$n_- = -1, \tag{6.10}$$

$$n_a = \omega\bar{\xi}_a + \bar{\omega}\xi_a, \tag{6.11}$$

where we used the equation (5.11) and (5.16) in section V.

Let us now write the main equations, the trivial equation, and the supplementary equations in the Newman-Unti coordinate. $G^{AB} l_B$ can be written as

$$G^{AB} l_B = -G^{AB}\delta_B^{\ +} = -G^{+A}. \tag{6.12}$$

Therefore we have

$$G^{AB} l_B = 0 \Longleftrightarrow \Big\{G^{++} = 0, \quad G^{+-} = 0, \quad G^{+a} = 0\Big\}. \tag{6.13}$$

Also $G^{AB} m_A m_B$ becomes

$$\begin{aligned} G^{AB} m_A m_B &= G^{++} m_+ m_+ + 2G^{+a} m_+ m_a + G^{ab} m_a m_b \\ &= G^{ab} m_a m_b \\ &= G^{ab} \xi_a \xi_b, \end{aligned} \tag{6.14}$$

where we used the equation (6.13). Thus the main equations (6.1) are equivalent to the following set of equations

$$\left\{ \begin{array}{l} G^{AB}l_B = 0 \\ G^{AB}m_A m_B = 0 \end{array} \right\} \Longleftrightarrow \left\{ \begin{array}{lll} G^{++} = 0, & G^{+-} = 0, & G^{+a} = 0 \\ G^{ab}\xi_a \xi_b = 0 & & \end{array} \right\}. \tag{6.15}$$

Notice that $G^{AB}m_A\bar{m}_B$ becomes,

$$\begin{aligned} G^{AB}m_A\bar{m}_B &= G^{++}m_+\bar{m}_+ + G^{+a}m_+\bar{m}_a + G^{a+}m_a\bar{m}_+ + G^{ab}m_a\bar{m}_b \\ &= G^{ab}m_{(a}\bar{m}_{b)} \\ &= G^{ab}\xi_{(a}\bar{\xi}_{b)}, \end{aligned} \tag{6.16}$$

using the main equations. Therefore we have the following correspondence

$$G^{AB}m_A\bar{m}_B = 0 \Longleftrightarrow G^{ab}\xi_{(a}\bar{\xi}_{b)} = 0, \tag{6.17}$$

assuming that the main equations hold. Let us consider the supplementary equations now. Using the main equations (6.15), $G^{AB}n_A m_B$ becomes

$$\begin{aligned} G^{AB}n_A m_B &= G^{++}n_+m_+ + G^{+a}n_+m_a + G^{-+}n_-m_+ + G^{-a}n_-m_a + G^{a+}n_a m_+ \\ &\quad + G^{ab}n_a m_b \\ &= G^{-a}n_-m_a + G^{ab}n_a m_b \\ &= -G^{-a}\xi_a + \bar{\omega}G^{ab}\xi_a\xi_b + \omega G^{ab}\bar{\xi}_{(a}\xi_{b)} \\ &= -G^{-a}\xi_a, \end{aligned} \tag{6.18}$$

where in the last line we used the equations

$$G^{ab}\xi_a\xi_b = 0 = G^{ab}\bar{\xi}_{(a}\xi_{b)}, \tag{6.19}$$

which follow from the equations (6.15) and (6.17). Thus, provided that the main equations and the trivial equation hold, we find that

$$G^{AB}n_A m_B = 0 = -G^{-a}\xi_a. \tag{6.20}$$

This equation, together with its complex conjugate

$$G^{-a}\bar{\xi}_a = 0, \tag{6.21}$$

yields

$$2G^{-a}\xi_{(a}\bar{\xi}_{b)} = G^{-a}\phi_{ab} = 0. \tag{6.22}$$

This shows that the following sets of equations are equivalent,

$$G^{AB}n_A m_B = 0 \Longleftrightarrow G^{-a} = 0, \tag{6.23}$$

provided the main equations and the trivial equation hold. Finally let us notice that $G^{AB}n_A n_B$ becomes

$$
\begin{aligned}
G^{AB}n_A n_B &= G^{++}n_+n_+ + G^{--}n_-n_- + G^{ab}n_a n_b + 2G^{+-}n_+n_- + 2G^{+a}n_+n_a \\
&\quad +2G^{-a}n_-n_a \\
&= G^{--}n_-n_- + G^{ab}n_a n_b + 2G^{-a}n_-n_a \\
&= G^{--} + G^{ab}\Big(\omega\bar{\xi}_a + \bar{\omega}\xi_a\Big)\Big(\omega\bar{\xi}_b + \bar{\omega}\xi_b\Big) - 2G^{-a}\Big(\omega\bar{\xi}_a + \bar{\omega}\xi_a\Big) \\
&= G^{--}, \qquad (6.24)
\end{aligned}
$$

where we again used the main equations and the trivial equation. From (6.23) and (6.24), we find that the followings are equivalent:

$$
\left\{ \begin{array}{l} G^{AB}n_A m_B = 0 \\ G^{AB}n_A n_B = 0 \end{array} \right\} \Longleftrightarrow \left\{ \begin{array}{l} G^{-a} = 0 \\ G^{--} = 0 \end{array} \right\}. \qquad (6.25)
$$

Putting these results together, the Einstein's equations in the standard null hypersurface formalism become, in the Newman-Unti coordinate, as follows:

$$
\begin{aligned}
&(A)\ \text{main equations}: \left\{ \begin{array}{l} G^{AB}l_B = 0 \\ G^{AB}m_A m_B = 0 \end{array} \right\} \Longleftrightarrow \left\{ \begin{array}{l} G^{++} = 0,\ G^{+-} = 0,\ G^{+a} = 0 \\ G^{ab}\xi_a\xi_b = 0 \end{array} \right\}, \\
&(B)\ \text{trivial equation}: G^{AB}m_A\bar{m}_B = 0 \Longleftrightarrow G^{ab}\xi_{(a}\bar{\xi}_{b)} = 0, \qquad (6.26) \\
&(C)\ \text{supplementary equations}: \left\{ \begin{array}{l} G^{AB}n_A m_B = 0 \\ G^{AB}n_A n_B = 0 \end{array} \right\} \Longleftrightarrow \left\{ \begin{array}{l} G^{-a} = 0 \\ G^{--} = 0 \end{array} \right\}.
\end{aligned}
$$

Having expressed the main equations, trivial equation, and supplementary equations in the Newman-Unti coordinate, we now wish to group the KK field equations (4.2), $\cdots$, (4.8) in accordance with the above classification. Let S be the Einstein-Hilbert action. The variation of S with respect to the metric[14] g_{AB} is given by

$$
\begin{aligned}
\delta S &= \int G^{AB}\delta g_{AB} + \text{surface terms} \\
&= \int G^{\mu\nu}\delta g_{\mu\nu} + \int 2G^{\mu a}\delta g_{\mu a} + \int G^{ab}\delta g_{ab} + \text{surface terms}, \qquad (6.27)
\end{aligned}
$$

where the integration is over 4-dimensional spacetime. In terms of the KK variables (2.2) in section II, the metric variations can be written as

$$
\begin{aligned}
\delta g_{\mu\nu} &= \delta\gamma_{\mu\nu} + A_\mu{}^a A_\nu{}^b\delta\phi_{ab} + \phi_{ab}A_\mu{}^a\delta A_\nu{}^b + \phi_{ab}A_\nu{}^b\delta A_\mu{}^a, \qquad (6.28) \\
\delta g_{\mu a} &= A_\mu{}^b\delta\phi_{ab} + \phi_{ab}\delta A_\mu{}^b, \qquad (6.29) \\
\delta g_{ab} &= \delta\phi_{ab}. \qquad (6.30)
\end{aligned}
$$

Therefore δS becomes

$$
\begin{aligned}
\delta S &= \int G^{\mu\nu}\delta\gamma_{\mu\nu} + \int 2\Big(G^{\mu\nu}\phi_{ab}A_\nu{}^b + G^{\mu b}\phi_{ab}\Big)\delta A_\mu{}^a \\
&\quad + \int \Big(G^{\mu\nu}A_\mu{}^a A_\nu{}^b + 2G^{\mu a}A_\mu{}^b + G^{ab}\Big)\delta\phi_{ab} + \text{surface terms}. \qquad (6.31)
\end{aligned}
$$

---

[14] Here the notations are as before, i.e. $A = (\mu, a); \mu = (0,1)$ or $(+,-), a = (2,3)$.

If we equate the coefficients of each variation in (6.31) to zero, we could obtain the Einstein's equations written in terms of the general KK variables without any gauge-fixing. If we use the gauge condition (3.2) in section III at the end of the above variations, then we find that[15]

$$\begin{aligned}\delta S = \int \Big\{ & G_{\rm p}^{++}\delta\gamma_{++} + 2G_{\rm p}^{+-}\delta\gamma_{+-} + G_{\rm p}^{--}\delta\gamma_{--} + 2\Big(G_{\rm p}^{++}\phi_{ab}A_+{}^b + G_{\rm p}^{+-}\phi_{ab}A_-{}^b + G_{\rm p}^{+b}\phi_{ab}\Big)\delta A_+{}^a \\ & +2\Big(G_{\rm p}^{-+}\phi_{ab}A_+{}^b + G_{\rm p}^{--}\phi_{ab}A_-{}^b + G_{\rm p}^{-b}\phi_{ab}\Big)\delta A_-{}^a \\ & +\Big(G_{\rm p}^{++}A_+{}^a A_+{}^b + 2G_{\rm p}^{+-}A_+{}^a A_-{}^b + G_{\rm p}^{--}A_-{}^a A_-{}^b + 2G_{\rm p}^{+a}A_+{}^b + 2G_{\rm p}^{-a}A_-{}^b + G_{\rm p}^{ab}\Big)\delta\phi_{ab}\Big\} \\ & +\text{surface terms.} \end{aligned} \qquad (6.32)$$

The equations resulting from these variations will be precisely the KK field equations, i.e. the equations (4.2), $\cdots$, (4.8) in section IV. Notice that the equations of motion of ϕ_{ab},

$$G_{\rm p}^{++}A_+{}^a A_+{}^b + 2G_{\rm p}^{+-}A_+{}^a A_-{}^b + G_{\rm p}^{--}A_-{}^a A_-{}^b + 2G_{\rm p}^{+a}A_+{}^b + 2G_{\rm p}^{-a}A_-{}^b + G_{\rm p}^{ab} = 0, \qquad (6.33)$$

can be split into three equations by taking trace with ϕ_{ab}, and by multiplying with $\xi_a\xi_b$ and its complex conjugate. The resultant equations are

$$\begin{aligned} (d)\quad & G_{\rm p}^{++}\phi_{ab}A_+{}^a A_+{}^b + 2G_{\rm p}^{+-}\phi_{ab}A_+{}^a A_-{}^b + G_{\rm p}^{--}\phi_{ab}A_-{}^a A_-{}^b + 2G_{\rm p}^{+a}\phi_{ab}A_+{}^b \\ & +2G_{\rm p}^{-a}\phi_{ab}A_-{}^b + G_{\rm p}^{ab}\phi_{ab} = 0, & (6.34) \\ (e)\quad & G_{\rm p}^{++}(\xi_a A_+{}^a)^2 + 2G_{\rm p}^{+-}(\xi_a A_+{}^a)(\xi_b A_-{}^b) + G_{\rm p}^{--}(\xi_a A_-{}^a)^2 + 2G_{\rm p}^{+a}\xi_a\xi_b A_+{}^b \\ & +2G_{\rm p}^{-a}\xi_a\xi_b A_-{}^b + G_{\rm p}^{ab}\xi_a\xi_b = 0, \ \text{ and c.c.,} & (6.35) \end{aligned}$$

which correspond to the equations (4.5) and (4.6) in section IV obtained by the variations with respect to σ and ρ_{ab} ($\det\rho_{ab} = 1$). Thus, together with the equations (6.34) and (6.35), the complete set of the KK field equations in terms of the Einstein tensor becomes, in the Polyakov gauge,

$$\begin{aligned} (a)\quad & G_{\rm p}^{++} = 0, & (6.36) \\ (b)\quad & G_{\rm p}^{++}\phi_{ab}A_+{}^b + G_{\rm p}^{+-}\phi_{ab}A_-{}^b + G_{\rm p}^{+b}\phi_{ab} = 0, & (6.37) \\ (c)\quad & G_{\rm p}^{-+}\phi_{ab}A_+{}^b + G_{\rm p}^{--}\phi_{ab}A_-{}^b + G_{\rm p}^{-b}\phi_{ab} = 0, & (6.38) \\ (f)\quad & G_{\rm p}^{+-} = 0, & (6.39) \\ (g)\quad & G_{\rm p}^{--} = 0. & (6.40) \end{aligned}$$

Notice that the right hand side of the classification (6.26) requires the main equations be valid throughout. Thus, if we assume that $G_{\rm p}^{++} = 0$, $G_{\rm p}^{+-} = 0$, $G_{\rm p}^{+a} = 0$, and $A_-{}^a = 0$, the equations (6.34), $\cdots$, (6.40) become,

---

[15] Here the subscript $_{\rm p}$ in $G_{\rm p}^{AB}$ stands for the Einstein tensor G^{AB} evaluated in the Polyakov gauge (3.2) in section III.

$$(a) \quad G_{\rm p}^{++} = 0, \tag{6.41}$$
$$(b) \quad G_{\rm p}^{+a} = 0, \tag{6.42}$$
$$(c) \quad G_{\rm p}^{-a} = 0, \tag{6.43}$$
$$(d) \quad G_{\rm p}^{ab}\xi_{(a}\bar{\xi}_{b)} = 0, \tag{6.44}$$
$$(e) \quad G_{\rm p}^{ab}\xi_{(a}\xi_{b)} = 0, \ \ \text{and c.c.}, \tag{6.45}$$
$$(f) \quad G_{\rm p}^{+-} = 0, \tag{6.46}$$
$$(g) \quad G_{\rm p}^{--} = 0. \tag{6.47}$$

It is well-known that the four equations $G^{-A} = 0$, when $x^{-} = v$ is viewed as the time coordinate, are the four constraint equations of the vacuum general relativity. These are the equations (6.43), (6.46), and (6.47), which correspond in the KK formalism to the constraint equations (4.4), (4.7), and (4.8) in section IV, since only the first v-time derivatives appear in these equations.

VII. DISCUSSIONS

In this paper, we presented the KK formalism of general relativity of 4-dimensional spacetimes, viewing the spacetime as a fibred manifold, namely, a local product of the (1+1)-dimensional base manifold and the 2-dimensional fibre space. Within this framework, we made a decomposition of a given 4-dimensional spacetime metric into sets of fields which transform as a tensor field, gauge fields, and scalar fields, with respect to the group of diffeomorphisms of the 2-dimensional fibre space. Moreover, using the Polyakov gauge, we arrived at an amazingly simple expression of the Einstein-Hilbert action, i.e. a Yang-Mills type gauge theory action coupled to a non-linear sigma field and a scalar field. Then we presented the Einstein's equations in terms of these fields.

This KK approach has several advantages, as we already mentioned. In connection with issues of quantum gravity, this approach has the following aspects which deserve further remarks. First, the problem of solving the Einstein's constraint equations is known to be one of the most important tasks in quantum general relativity. Remarkably, in the (2,2)-KK approach, the diffeomorphisms of the 2-dimensional fibre space turns out to be a local gauge symmetry, exactly as in Yang-Mills theory. Therefore the two constraint equations associated with these diffeomorphisms can be automatically solved, by using the diffN_2-invariant quantities. However, there are two additional constraint equations which require further studies [5] in order to fully take care of the four Einstein's constraint equations.

Second, the Polyakov gauge is shown to be equivalent to the Newman-Unti coordinate, modulo the diffN_2 transformations. It is well-known that, in the null hypersurface formalism in the Newman-Unti coordinate, the physical gravitational degrees of freedom can be easily isolated, whereas in the standard (3+1)-formalism or in the ADM formalism such an isolation can be done only in the linearized theory. Hence the KK formalism in the Polyakov gauge shares this salient feature of the null hypersurface formalism: it is the non-linear sigma field ρ_{ab} with a unit determinant that is precisely the physical gravitational degrees of freedom, which are also known as the conformal 2-metric of the transverse 2-surface in the null hypersurface formalism.

Third, it is also of interest to find how the lightcones fluctuate in quantum gravity. In the formalism it is quite easy to address this issue. Recall that a lightcone is a collection of null vector fields originating from a given spacetime point. The relevant null vector field is, as given in section V (with $\omega = 0$),

$$n = \frac{\partial}{\partial u} - h\frac{\partial}{\partial v} - A_{+}{}^{a}\frac{\partial}{\partial y^{a}}, \tag{7.1}$$

which clearly has a zero norm. This vector field is controlled by the fields h and $A_{+}{}^{a}$. Therefore, by quantizing these fields, we can quantize the lightcone itself without referring to any background spacetime geometry.

Fourth, it should be mentioned that the Lie derivative appears naturally in this formalism, via the *minimal* couplings to the diffN_2-valued gauge fields. In the standard (3+1)-formalism, the natural derivative operator is the metric-compatible covariant derivative, which requires the metric be non-degenerate. The Lie derivative, on the other hand, can be defined even when the metric is degenerate. Therefore, the KK formalism, based on the notion of the Lie derivative, should be extendable to spacetimes where the metric is degenerate, which would be hard to describe in conventional approaches.

One may also attack more "practical" problems in classical general relativity. For instance, we expect the KK formalism to be useful for problems such as finding exact solutions of the Einstein's equations [24], in view of the fact that the null hypersurface formalism, particularly the Newman-Penrose version, has been proven extremely useful for such problems. To those problems concerned with asymptotically flat radiative spacetimes [5] or gravitational waves in general, the new formalism may be also applicable. A related but probably more challenging problem is to reinterpret *all* the known exact solutions of the Einstein's equations from the perspective of the Yang-Mills type gauge theories coupled to the non-linear sigma field and a scalar field on the (1+1)-dimensional base manifold. This seems very interesting, for there are a number of exact solutions of the Einstein's equations which do not permit a sensible physical interpretation from the 4-dimensional spacetime perspective [25].

Finally, we would like to remark on the question of finiteness of quantum gravity. Since the question whether a theory is renormalizable or not depends critically on what spacetime dimensions the theory is defined on, this (1+1)-dimensional description of general relativity of 4-dimensions may lead to a happy conclusion that quantum gravity is, after all, a *finite* theory, at least when viewed from this (1+1)-dimensional field theory perspective.

Acknowledgments

This work is supported in part by the Basic Science Research Institute Program, Ministry of Education, 1995 (BSRI-95-2418), and by non-directed research fund, Korea Research Foundation, 1995 (04-D-0073). It is also supported by Korea Science and Engineering Foundation (95-0702-04-01-3), and by the SRC program of SNU-CTP of KOSEF. Parts of this work were done while the author was a post-doctoral fellow at FAMAF, the Universidad Nacional de Córdoba, Argentina.

APPENDIX A: DIFFN_2 TRANSFORMATIONS

In this Appendix we shall first derive the finite transformation properties (2.5), (2.6), and (2.7) and the corresponding infinitesimal transformation properties (2.9), (2.10), and (2.11), and then show that $D_\mu\phi_{ab}$ and $F_{\mu\nu}{}^a$ transform covariantly under the diffN_2 transformations, as given in (2.16) and (2.17) in section II. The spacetime metric at the point (x^μ, y^a) is given by (2.2) in section II,

$$\begin{aligned} ds^2 &= \gamma_{\mu\nu}dx^\mu dx^\nu + \phi_{ab}\Big(dy^a + A_\mu{}^a dx^\mu\Big)\Big(dy^b + A_\nu{}^b dx^\nu\Big) \\ &= \Big(\gamma_{\mu\nu} + \phi_{ab}A_\mu{}^a A_\nu{}^b\Big)dx^\mu dx^\nu + 2\phi_{ab}A_\mu{}^a dx^\mu dy^b + \phi_{ab}dy^a dy^b. \end{aligned} \tag{A1}$$

Let us consider the following arbitrary transformation of the y^a-coordinate at each point of the base manifold x^μ, while keeping x^μ constant,

$$\begin{aligned} y^{'a} &= y^{'a}(x,y), \\ x^{'\mu} &= x^\mu, \end{aligned} \tag{A2}$$

where $y^{'a}(x,y)$ is an arbitrary function of x^μ and y^a. Then we have

$$\begin{aligned} dy^{'a} &= \Big(\frac{\partial y^{'a}}{\partial y^c}\Big)dy^c + \Big(\frac{\partial y^{'a}}{\partial x^\mu}\Big)dx^\mu, \\ dx^{'\mu} &= dx^\mu, \end{aligned} \tag{A3}$$

or equivalently,

$$\begin{aligned} dy^a &= \frac{\partial y^a}{\partial y^{'c}}\Big\{dy^{'c} - \Big(\frac{\partial y^{'c}}{\partial x^\mu}\Big)dx^{'\mu}\Big\}, \\ dx^\mu &= dx^{'\mu}. \end{aligned} \tag{A4}$$

Thus we can express the metric at the point (x^μ, y^a) in terms of the differentials of the new coordinates $dx^{'\mu}$ and $dy^{'a}$. The term proportional to $dx^\mu dy^a$ in (A1) becomes

$$\begin{aligned} &2\phi_{ab}(x,y)A_\mu{}^a(x,y)dx^\mu dy^b \\ &= 2\phi_{ab}(x,y)A_\mu{}^a(x,y)dx^{'\mu}\Big(\frac{\partial y^b}{\partial y^{'c}}\Big)\Big\{dy^{'c} - \Big(\frac{\partial y^{'c}}{\partial x^\nu}\Big)dx^{'\nu}\Big\} \\ &= 2\Big(\frac{\partial y^b}{\partial y^{'c}}\Big)\phi_{ab}(x,y)\Big(\frac{\partial y^a}{\partial y^{'d}}\Big)\Big(\frac{\partial y^{'d}}{\partial y^e}\Big)A_\mu{}^e(x,y)dx^{'\mu}\Big\{dy^{'c} - \Big(\frac{\partial y^{'c}}{\partial x^\nu}\Big)dx^{'\nu}\Big\} \\ &= 2\Big(\frac{\partial y^a}{\partial y^{'c}}\Big)\Big(\frac{\partial y^b}{\partial y^{'d}}\Big)\phi_{ab}(x,y)\Big(\frac{\partial y^{'d}}{\partial y^e}\Big)A_\mu{}^e(x,y)dx^{'\mu}\Big\{dy^{'c} - \Big(\frac{\partial y^{'c}}{\partial x^\nu}\Big)dx^{'\nu}\Big\}, \end{aligned} \tag{A5}$$

where the (x^μ, y^a)-dependence of each field is shown explicitly, and the identity

$$\Big(\frac{\partial y^a}{\partial y^{'d}}\Big)\Big(\frac{\partial y^{'d}}{\partial y^e}\Big) = \delta_e{}^a \tag{A6}$$

was used. The term containing $dy^a dy^b$ becomes

$$\begin{aligned}
&\phi_{ab}(x,y)dy^a dy^b \\
&= \phi_{ab}(x,y)\Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\Big\{dy'^c - \Big(\frac{\partial y'^c}{\partial x^\mu}\Big)dx'^\mu\Big\}\Big\{dy'^d - \Big(\frac{\partial y'^d}{\partial x^\nu}\Big)dx'^\nu\Big\} \\
&= \Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\phi_{ab}(x,y)\Big\{dy'^c dy'^d - 2\Big(\frac{\partial y'^d}{\partial x^\mu}\Big)dy'^c dx'^\mu + \Big(\frac{\partial y'^c}{\partial x^\mu}\Big)\Big(\frac{\partial y'^d}{\partial x^\nu}\Big)dx'^\mu dx'^\nu\Big\}.
\end{aligned} \tag{A7}$$

Therefore the line element (A1) becomes

$$\begin{aligned}
ds^2 &= \Big\{\gamma_{\mu\nu}(x,y) + \phi_{ab}(x,y)A_\mu{}^a(x,y)A_\nu{}^b(x,y)\Big\}dx'^\mu dx'^\nu \\
&\quad + \Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\phi_{ab}(x,y)\Big\{dy'^c dy'^d - 2\Big(\frac{\partial y'^d}{\partial x^\mu}\Big)dy'^c dx'^\mu + \Big(\frac{\partial y'^c}{\partial x^\mu}\Big)\Big(\frac{\partial y'^d}{\partial x^\nu}\Big)dx'^\mu dx'^\nu\Big\} \\
&\quad + 2\Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\phi_{ab}(x,y)\Big(\frac{\partial y'^d}{\partial y^e}\Big)A_\mu{}^e(x,y)dx'^\mu\Big\{dy'^c - \Big(\frac{\partial y'^c}{\partial x^\nu}\Big)dx'^\nu\Big\} \\
&= \gamma_{\mu\nu}(x,y)dx'^\mu dx'^\nu + \Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\phi_{ab}(x,y)dy'^c dy'^d \\
&\quad + 2\Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\phi_{ab}(x,y)\Big\{\Big(\frac{\partial y'^d}{\partial y^e}\Big)A_\mu{}^e(x,y) - \frac{\partial y'^d}{\partial x^\mu}\Big\}dx'^\mu dy'^c \\
&\quad + \phi_{ab}(x,y)\Big\{A_\mu{}^a(x,y)A_\nu{}^b(x,y) - 2\Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\Big(\frac{\partial y'^d}{\partial y^e}\Big)A_\mu{}^e(x,y)\Big(\frac{\partial y'^c}{\partial x^\nu}\Big) \\
&\quad + \Big(\frac{\partial y^a}{\partial y'^c}\Big)\Big(\frac{\partial y^b}{\partial y'^d}\Big)\Big(\frac{\partial y'^c}{\partial x^\mu}\Big)\Big(\frac{\partial y'^d}{\partial x^\nu}\Big)\Big\}dx'^\mu dx'^\nu,
\end{aligned} \tag{A8}$$

which must be equal to

$$ds'^2 = \gamma'_{\mu\nu}(x',y')dx'^\mu dx'^\nu + \phi'_{ab}(x',y')\Big\{dy'^a + A'_\mu{}^a(x',y')dx'^\mu\Big\}\Big\{dy'^b + A'_\nu{}^b(x',y')dx'^\nu\Big\}, \tag{A9}$$

due to the diffeomorphism invariance. If we compare terms containing $dy'^a dy'^b$ in (A8) and (A9), we find that

$$\phi'_{ab}(x',y') = \Big(\frac{\partial y^c}{\partial y'^a}\Big)\Big(\frac{\partial y^d}{\partial y'^b}\Big)\phi_{cd}(x,y), \tag{A10}$$

which shows that $\phi_{cd}(x,y)$ is a tensor field with respect to the diffN_2 transformations. Then we may write the line element (A8) as

$$\begin{aligned}
ds^2 &= \gamma_{\mu\nu}(x,y)dx'^\mu dx'^\nu + \phi'_{cd}(x',y')dy'^c dy'^d + 2\phi'_{cd}(x',y')\Big\{\Big(\frac{\partial y'^d}{\partial y^a}\Big)A_\mu{}^a(x,y) - \frac{\partial y'^d}{\partial x^\mu}\Big\}dx'^\mu dy'^c \\
&\quad + \phi'_{cd}(x',y')\Big\{\Big(\frac{\partial y'^c}{\partial y^a}\Big)A_\mu{}^a(x,y) - \frac{\partial y'^c}{\partial x^\mu}\Big\}\Big\{\Big(\frac{\partial y'^d}{\partial y^b}\Big)A_\nu{}^b(x,y) - \frac{\partial y'^d}{\partial x^\nu}\Big\}dx'^\mu dx'^\nu.
\end{aligned} \tag{A11}$$

From this expression, we find the transformation properties of $A_\mu{}^a(x,y)$ and $\gamma_{\mu\nu}(x,y)$,

$$A'_\mu{}^a(x',y') = \Big(\frac{\partial y'^a}{\partial y^b}\Big)A_\mu{}^b(x,y) - \frac{\partial y'^a}{\partial x^\mu}(x,y), \tag{A12}$$

$$\gamma'_{\mu\nu}(x',y') = \gamma_{\mu\nu}(x,y). \tag{A13}$$

The equations (A10), (A12), and (A13) are the equations (2.5), (2.6), and (2.7) in section II.

The infinitesimal versions of the above finite transformations can be also found by considering the following transformations

$$\begin{aligned} &{y'}^{a} = y^{a} + \xi^{a}(x,y) \qquad (\mathrm{O}(\xi^2) \ll 1), \\ &{x'}^{\mu} = x^{\mu}, \end{aligned} \tag{A14}$$

where ξ^a is an arbitrary, infinitesimal function of x^μ and y^a. Thus we have

$$\frac{\partial {y'}^{a}}{\partial y^{c}} = \delta_{c}^{\ a} + \frac{\partial \xi^{a}}{\partial y^{c}}, \tag{A15}$$

or

$$\frac{\partial y^{c}}{\partial {y'}^{a}} = \delta_{a}^{\ c} - \frac{\partial \xi^{c}}{\partial y^{a}} + \cdots, \tag{A16}$$

where $\cdots$ means terms of $\mathrm{O}(\xi^2)$. Let us expand the left hand side of the equation (A10) using (A14). It becomes

$$\phi'_{ab}(x', y+\xi) = \phi'_{ab}(x,y) + \xi^{c}\frac{\partial}{\partial y^{c}}\phi_{ab}(x,y) + \cdots, \tag{A17}$$

whereas the right hand side becomes

$$\begin{aligned} \Big(\frac{\partial y^{c}}{\partial {y'}^{a}}\Big)\Big(\frac{\partial y^{d}}{\partial {y'}^{b}}\Big)\phi_{cd}(x,y) &= \Big(\delta_{a}^{\ c} - \frac{\partial \xi^{c}}{\partial y^{a}} + \cdots\Big)\Big(\delta_{b}^{\ d} - \frac{\partial \xi^{d}}{\partial y^{b}} + \cdots\Big)\phi_{cd}(x,y) \\ &= \phi_{ab}(x,y) - \frac{\partial \xi^{c}}{\partial y^{a}}\phi_{cb}(x,y) - \frac{\partial \xi^{c}}{\partial y^{b}}\phi_{ac}(x,y) + \cdots. \end{aligned} \tag{A18}$$

From (A17) and (A18), we therefore have

$$\begin{aligned} \delta\phi_{ab}(x,y) &\equiv \phi'_{ab}(x,y) - \phi_{ab}(x,y) \\ &= -\xi^{c}\frac{\partial}{\partial y^{c}}\phi_{ab}(x,y) - \frac{\partial \xi^{c}}{\partial y^{a}}\phi_{cb}(x,y) - \frac{\partial \xi^{c}}{\partial y^{b}}\phi_{ac}(x,y) \\ &= -[\xi, \phi]_{\mathrm{L}ab}, \end{aligned} \tag{A19}$$

where the subscript $_{\mathrm{L}}$ denotes the Lie derivative of ϕ_{ab} along the vector field $\xi = \xi^{c}\partial/\partial y^{a}$, i.e.

$$[\xi, \phi]_{\mathrm{L}ab} = \xi^{c}\frac{\partial}{\partial y^{c}}\phi_{ab} + \frac{\partial \xi^{c}}{\partial y^{a}}\phi_{cb} + \frac{\partial \xi^{c}}{\partial y^{b}}\phi_{ac}. \tag{A20}$$

It is a straightforward exercise to derive the infinitesimal transformation properties $A_{\mu}^{\ a}$ and $\gamma_{\mu\nu}$ from the finite transformations (A12) and (A13). They are found to be

$$\begin{aligned} \delta A_{\mu}^{\ a}(x,y) &= -\frac{\partial \xi^{a}}{\partial x^{\mu}} + [A_{\mu},\ \xi]_{\mathrm{L}}^{a} \\ &= -D_{\mu}\xi^{a}, \end{aligned} \tag{A21}$$

$$\delta\gamma_{\mu\nu}(x,y) = -[\xi, \gamma_{\mu\nu}]_{\mathrm{L}}, \tag{A22}$$

where the Lie brackets are defined as

$$
\begin{aligned}
&[A_\mu,\ \xi]^a_{\rm L} = A_\mu{}^c \frac{\partial \xi^a}{\partial y^c} - \xi^c \frac{\partial}{\partial y^c} A_\mu{}^a, \\
&[\xi, \gamma_{\mu\nu}]_{\rm L} = \xi^a \frac{\partial}{\partial y^a} \gamma_{\mu\nu}.
\end{aligned} \tag{A23}
$$

The equations (A19), (A21), and (A22) show that ϕ_{ab}, $A_\mu{}^a$, and $\gamma_{\mu\nu}$ behave like a tensor field, gauge fields, and scalar fields under the diffN_2 transformation, respectively. These are the equations (2.9),(2.10), and (2.11) in section II.

Let us also derive the following transformation properties of $D_\mu \phi_{ab}$ and $F_{\mu\nu}{}^a$ under the diffN_2 transformations,

$$
\delta(D_\mu \phi_{ab}) = -[\xi, D_\mu \phi]_{{\rm L}ab}, \tag{A24}
$$

$$
\delta F_{\mu\nu}{}^a = -[\xi, F_{\mu\nu}]^a_{\rm L}, \tag{A25}
$$

as given in equations (2.16) and (2.17) in section II. Notice that

$$
\begin{aligned}
\delta(D_\mu \phi_{ab}) &= \partial_\mu(\delta\phi_{ab}) - (\delta A_\mu{}^c)(\partial_c \phi_{ab}) - A_\mu{}^c \partial_c(\delta \phi_{ab}) - \partial_a(\delta A_\mu{}^c)\phi_{bc} \\
&\quad -(\partial_a A_\mu{}^c)\delta\phi_{bc} - \partial_b(\delta A_\mu{}^c)\phi_{ac} - (\partial_b A_\mu{}^c)\delta\phi_{ac} \\
&= -\partial_\mu\Big([\xi, \phi]_{{\rm L}ab}\Big) + (D_\mu \xi^c)(\partial_c \phi_{ab}) + A_\mu{}^c \partial_c\Big([\xi, \phi]_{{\rm L}ab}\Big) + \partial_a(D_\mu \xi^c)\phi_{bc} \\
&\quad +(\partial_a A_\mu{}^c)[\xi, \phi]_{{\rm L}bc} + \partial_b(D_\mu \xi^c)\phi_{ac} + (\partial_b A_\mu{}^c)[\xi, \phi]_{{\rm L}ac} \\
&= -\partial_\mu\Big([\xi, \phi]_{{\rm L}ab}\Big) + [A_\mu, [\xi, \phi]_{\rm L}]_{{\rm L}ab} + [D_\mu \xi, \phi]_{{\rm L}ab},
\end{aligned} \tag{A26}
$$

using the equations (A19) and (A21), and the following Lie brackets

$$
[A_\mu, [\xi, \phi]_{\rm L}]_{{\rm L}ab} = A_\mu{}^c \partial_c\Big([\xi, \phi]_{{\rm L}ab}\Big) + (\partial_a A_\mu{}^c)[\xi, \phi]_{{\rm L}bc} + (\partial_b A_\mu{}^c)[\xi, \phi]_{{\rm L}ac}, \tag{A27}
$$

$$
[D_\mu \xi, \phi]_{{\rm L}ab} = (D_\mu \xi^c)(\partial_c \phi_{ab}) + \partial_a(D_\mu \xi^c)\phi_{bc} + \partial_b(D_\mu \xi^c)\phi_{ac}. \tag{A28}
$$

Using the Leibniz rule of the derivative ∂_μ

$$
\partial_\mu\Big([\xi, \phi]_{{\rm L}ab}\Big) = [\partial_\mu \xi, \phi]_{{\rm L}ab} + [\xi, \partial_\mu \phi]_{{\rm L}ab}, \tag{A29}
$$

and the properties of the Lie bracket

$$
[D_\mu \xi, \phi]_{{\rm L}ab} = [\partial_\mu \xi, \phi]_{{\rm L}ab} - [[A_\mu, \xi]_{\rm L}, \phi]_{{\rm L}ab}, \tag{A30}
$$

$$
[A_\mu, [\xi, \phi]_{\rm L}]_{{\rm L}ab} = -[\xi, [\phi, A_\mu]_{\rm L}]_{{\rm L}ab} - [\phi, [A_\mu, \xi]_{\rm L}]_{{\rm L}ab}, \tag{A31}
$$

we find that

$$
\begin{aligned}
\delta(D_\mu \phi_{ab}) &= -[\xi, \partial_\mu \phi]_{{\rm L}ab} + [\xi, [A_\mu, \phi]_{\rm L}]_{{\rm L}ab} \\
&= -[\xi, D_\mu \phi]_{{\rm L}ab}.
\end{aligned} \tag{A32}
$$

The infinitesimal variation of $F_{\mu\nu}{}^a$ can be obtained in a similar way. It becomes

$$
\begin{aligned}
\delta F_{\mu\nu}{}^{a} &= \partial_\mu(\delta A_\nu{}^a) - [\delta A_\mu, A_\nu]^a_{\mathrm{L}} - (\mu \leftrightarrow \nu) \\
&= -\partial_\mu\Big\{\partial_\nu \xi^a - [A_\nu, \xi]^a_{\mathrm{L}}\Big\} + [D_\mu \xi, A_\nu]^a_{\mathrm{L}} - (\mu \leftrightarrow \nu) \\
&= \partial_\mu\Big([A_\nu, \xi]^a_{\mathrm{L}}\Big) + [D_\mu \xi, A_\nu]^a_{\mathrm{L}} - (\mu \leftrightarrow \nu).
\end{aligned} \tag{A33}
$$

Using the following identities

$$\partial_\mu\Big([A_\nu, \xi]^a_{\mathrm{L}}\Big) = [\partial_\mu A_\nu, \xi]^a_{\mathrm{L}} + [A_\nu, \partial_\mu \xi]^a_{\mathrm{L}}, \tag{A34}$$

$$[D_\mu \xi, A_\nu]^a_{\mathrm{L}} = -[A_\nu, D_\mu \xi]^a_{\mathrm{L}} = -[A_\nu, \partial_\mu \xi]^a_{\mathrm{L}} + [A_\nu, [A_\mu, \xi]_{\mathrm{L}}]^a_{\mathrm{L}}, \tag{A35}$$

we find that

$$
\begin{aligned}
\delta F_{\mu\nu}{}^{a} &= [\partial_\mu A_\nu, \xi]^a_{\mathrm{L}} + [A_\nu, [A_\mu, \xi]_{\mathrm{L}}]^a_{\mathrm{L}} - (\mu \leftrightarrow \nu) \\
&= [\partial_\mu A_\nu - \partial_\nu A_\mu, \xi]^a_{\mathrm{L}} + [A_\nu, [A_\mu, \xi]_{\mathrm{L}}]^a_{\mathrm{L}} - [A_\mu, [A_\nu, \xi]_{\mathrm{L}}]^a_{\mathrm{L}} \\
&= -[\xi, \partial_\mu A_\nu - \partial_\nu A_\mu - [A_\mu, A_\nu]_{\mathrm{L}}]^a_{\mathrm{L}} \\
&= -[\xi, F_{\mu\nu}]^a_{\mathrm{L}},
\end{aligned} \tag{A36}
$$

where we used the Jacobi identity

$$[A_\nu, [A_\mu, \xi]_{\mathrm{L}}]^a_{\mathrm{L}} = -[A_\mu, [\xi, A_\nu]_{\mathrm{L}}]^a_{\mathrm{L}} - [\xi, [A_\nu, A_\mu]_{\mathrm{L}}]^a_{\mathrm{L}}. \tag{A37}$$

Therefore it follows that

$$\delta F_{\mu\nu}{}^{a} = -[\xi, F_{\mu\nu}]^a_{\mathrm{L}}. \tag{A38}$$

APPENDIX B: THE EINSTEIN-HILBERT ACTION IN THE (2,2)-SPLITTING

In this Appendix we shall describe the procedure of obtaining the scalar curvature (2.18) in section II of 4-dimensional spacetime. In the (2,2)-splitting of spacetime, we regard the 4-dimensional spacetime in question as a local product of a (1+1)-dimensional base manifold M_{1+1} and a 2-dimensional fibre space N_2, whose basis vector fields are ∂_μ and ∂_a, respectively. The (1+1)-dimensional space orthogonal to the fibre space N_2 is called the horizontal space, generated by the vector fields that can be written as linear combinations of ∂_μ and ∂_a,

$$\hat{\partial}_\mu = \partial_\mu - A_\mu{}^a \partial_a, \tag{B1}$$

where $A_\mu{}^a$ are the coefficient functions of (x^μ, y^a). Let $\hat{\partial}_a$ be such that

$$\hat{\partial}_a = \partial_a. \tag{B2}$$

The basis $(\hat{\partial}_\mu, \hat{\partial}_a)$ is called the horizontal lift basis [14], and the metric coefficients in this basis are given by

$$\hat{g}_{AB} = \begin{pmatrix} \gamma_{\mu\nu} & 0 \\ 0 & \phi_{ab} \end{pmatrix}. \tag{B3}$$

Here $\gamma_{\mu\nu}$ and ϕ_{ab} are the metrics of the horizontal space and the fibre space, respectively, and that

$$\hat{g}_{\mu a} = 0 \tag{B4}$$

is a consequence of the orthogonality of the horizontal space and the fibre space. In general, the horizontal lift basis vector fields are anholonomic,

$$[\hat{\partial}_A, \hat{\partial}_B] = f_{AB}{}^{C}\hat{\partial}_C, \tag{B5}$$

where $f_{AB}{}^{C}$ are the structure functions satisfying the relation

$$f_{AB}{}^{C} = f_{[AB]}{}^{C}. \tag{B6}$$

These structure functions are found to be

$$\begin{aligned} f_{\mu\nu}{}^{a} &= -f_{\nu\mu}{}^{a} = -F_{\mu\nu}{}^{a}, \\ f_{\mu a}{}^{b} &= -f_{a\mu}{}^{b} = \partial_a A_{\mu}{}^{b}, \end{aligned} \tag{B7}$$

and zero otherwise. Here $F_{\mu\nu}{}^{a}$ is defined as

$$F_{\mu\nu}{}^{a} = \partial_\mu A_{\nu}{}^{a} - \partial_\nu A_{\mu}{}^{a} - [A_\mu, A_\nu]^a_{\mathrm{L}}, \tag{B8}$$

as before.

The formulae of the connection coefficients and the curvature tensors in the anholonomic basis can be found in [14,15]. In the horizontal lift basis where the metric is given by (B3), they are given by

$$\hat{\Gamma}_{AB}{}^{C} = \frac{1}{2}\hat{g}^{CD}\Big(\hat{\partial}_A\hat{g}_{BD} + \hat{\partial}_B\hat{g}_{AD} - \hat{\partial}_D\hat{g}_{AB}\Big) + \frac{1}{2}\hat{g}^{CD}\Big(f_{ABD} - f_{BDA} - f_{ADB}\Big), \tag{B9}$$

$$\hat{R}_{ABC}{}^{D} = \hat{\partial}_A\hat{\Gamma}_{BC}{}^{D} - \hat{\partial}_B\hat{\Gamma}_{AC}{}^{D} + \hat{\Gamma}_{AE}{}^{D}\hat{\Gamma}_{BC}{}^{E} - \hat{\Gamma}_{BE}{}^{D}\hat{\Gamma}_{AC}{}^{E} - f_{AB}{}^{E}\hat{\Gamma}_{EC}{}^{D}, \tag{B10}$$

$$\hat{R}_{AC} = \hat{R}_{ABC}{}^{B}, \tag{B11}$$

$$R = \hat{g}^{AC}\hat{R}_{AC}, \tag{B12}$$

where $f_{ABC} = \hat{g}_{CD}f_{AB}{}^{D}$. Notice that in this basis $\hat{\Gamma}_{AB}{}^{C}$ is, in general, not symmetric:

$$\hat{\Gamma}_{AB}{}^{C} \neq \hat{\Gamma}_{BA}{}^{C} \quad \text{if} \quad f_{AB}{}^{C} \neq 0. \tag{B13}$$

In components the connection coefficients are given by

$$\hat{\Gamma}_{\mu\nu}{}^{\alpha} = \frac{1}{2}\gamma^{\alpha\beta}\Big(\hat{\partial}_\mu\gamma_{\nu\beta} + \hat{\partial}_\nu\gamma_{\mu\beta} - \hat{\partial}_\beta\gamma_{\mu\nu}\Big), \tag{B14}$$

$$\hat{\Gamma}_{\mu\nu}{}^{a} = -\frac{1}{2}\phi^{ab}\partial_b\gamma_{\mu\nu} - \frac{1}{2}F_{\mu\nu}{}^{a}, \tag{B15}$$

$$\hat{\Gamma}_{\mu a}{}^{\nu} = \hat{\Gamma}_{a\mu}{}^{\nu} = \frac{1}{2}\gamma^{\nu\alpha}\partial_a\gamma_{\mu\alpha} + \frac{1}{2}\gamma^{\nu\alpha}\phi_{ab}F_{\mu\alpha}{}^{b}, \tag{B16}$$

$$\hat{\Gamma}_{\mu a}{}^{b} = \frac{1}{2}\phi^{bc}\hat{\partial}_\mu\phi_{ac} + \frac{1}{2}\partial_a A_{\mu}{}^{b} - \frac{1}{2}\phi^{bc}\phi_{ae}\partial_c A_{\mu}{}^{e}$$

$$= \frac{1}{2}\phi^{bc} D_\mu \phi_{ac} + \partial_a A_\mu{}^b, \tag{B17}$$

$$\hat{\Gamma}_{a\mu}{}^b = \frac{1}{2}\phi^{bc}\hat{\partial}_\mu \phi_{ac} - \frac{1}{2}\partial_a A_\mu{}^b - \frac{1}{2}\phi^{bc}\phi_{ae}\partial_c A_\mu{}^e$$

$$= \frac{1}{2}\phi^{bc} D_\mu \phi_{ac}, \tag{B18}$$

$$\hat{\Gamma}_{ab}{}^\mu = -\frac{1}{2}\gamma^{\mu\nu}\hat{\partial}_\nu \phi_{ab} + \frac{1}{2}\gamma^{\mu\nu}\phi_{ac}\partial_b A_\nu{}^c + \frac{1}{2}\gamma^{\mu\nu}\phi_{bc}\partial_a A_\nu{}^c$$

$$= -\frac{1}{2}\gamma^{\mu\nu} D_\nu \phi_{ab}, \tag{B19}$$

$$\hat{\Gamma}_{ab}{}^c = \frac{1}{2}\phi^{cd}\Big(\partial_a \phi_{bd} + \partial_b \phi_{ad} - \partial_d \phi_{ab}\Big), \tag{B20}$$

where $\hat{\partial}_\mu \gamma_{\alpha\beta}$ and $\hat{\partial}_\mu \phi_{ab}$ are given by

$$\hat{\partial}_\mu \gamma_{\alpha\beta} = \partial_\mu \gamma_{\alpha\beta} - A_\mu{}^c \partial_c \gamma_{\alpha\beta}, \tag{B21}$$

$$\hat{\partial}_\mu \phi_{ab} = \partial_\mu \phi_{ab} - A_\mu{}^c \partial_c \phi_{ab}, \tag{B22}$$

and $D_\mu \phi_{ab}$ is defined as

$$D_\mu \phi_{ab} = \partial_\mu \phi_{ab} - A_\mu{}^c \partial_c \phi_{ab} - (\partial_a A_\mu{}^c)\phi_{bc} - (\partial_b A_\mu{}^c)\phi_{ac}$$

$$= \partial_\mu \phi_{ab} - [A_\mu, \phi]_{\mathrm{L}ab}. \tag{B23}$$

Here $[A_\mu, \phi]_{\mathrm{L}ab}$ is the Lie derivative of ϕ_{ab} along the vector field $A_\mu = A_\mu{}^a \partial_a$. From (B14), $\cdots$, (B20), we also find the following useful identities,

$$\hat{\Gamma}_{\mu\nu}{}^\mu = \hat{\Gamma}_{\nu\mu}{}^\mu = \frac{1}{2}\gamma^{\alpha\beta}\hat{\partial}_\nu \gamma_{\alpha\beta}, \tag{B24}$$

$$\hat{\Gamma}_{\mu a}{}^\mu = \hat{\Gamma}_{a\mu}{}^\mu = \frac{1}{2}\gamma^{\alpha\beta}\partial_a \gamma_{\alpha\beta}, \tag{B25}$$

$$\hat{\Gamma}_{\nu a}{}^a = \frac{1}{2}\phi^{ab} D_\nu \phi_{ab} + \partial_a A_\nu{}^a, \tag{B26}$$

$$\hat{\Gamma}_{a\nu}{}^a = \frac{1}{2}\phi^{ab} D_\nu \phi_{ab}, \tag{B27}$$

$$\hat{\Gamma}_{ab}{}^a = \frac{1}{2}\phi^{ac}\partial_b \phi_{ac}. \tag{B28}$$

The scalar curvature R of the 4-dimensional spacetime is given by

$$R = \hat{g}^{AB}\hat{R}_{AB}$$

$$= \gamma^{\mu\nu}\hat{R}_{\mu\nu} + \phi^{ac}\hat{R}_{ac}, \tag{B29}$$

where $\hat{R}_{\mu\nu}$ and $\hat{R}_{ac}$ are given by

$$\hat{R}_{\mu\nu} = \hat{R}_{\mu\alpha\nu}{}^\alpha + \hat{R}_{\mu a\nu}{}^a,$$

$$\hat{R}_{ac} = \hat{R}_{abc}{}^b + \hat{R}_{a\alpha c}{}^\alpha, \tag{B30}$$

respectively. The quantities $\gamma^{\mu\nu}\hat{R}_{\mu\nu}$ and $\phi^{ac}\hat{R}_{ac}$ are found to be

$$\begin{aligned}
\gamma^{\mu\nu}\hat{R}_{\mu\nu} &= \gamma^{\mu\nu}\tilde{R}_{\mu\nu} + \frac{1}{2}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b} + \frac{1}{4}\gamma^{\mu\nu}\phi^{ab}\phi^{cd}\Big\{(D_\mu\phi_{ac})(D_\nu\phi_{bd}) \\
&\qquad -(D_\mu\phi_{ab})(D_\nu\phi_{cd})\Big\} + \frac{1}{2}\Big(\hat{\nabla}_\mu + \hat{\Gamma}_{c\mu}{}^{c}\Big)j^\mu + \frac{1}{2}\Big(\hat{\nabla}_a + \hat{\Gamma}_{\alpha a}{}^{\alpha}\Big)j^a, \qquad \text{(B31)} \\
\phi^{ac}\hat{R}_{ac} &= \phi^{ac}R_{ac} - \frac{1}{4}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b} + \frac{1}{4}\phi^{ab}\gamma^{\mu\nu}\gamma^{\alpha\beta}\Big\{(\partial_a\gamma_{\mu\alpha})(\partial_b\gamma_{\nu\beta}) \\
&\qquad -(\partial_a\gamma_{\mu\nu})(\partial_b\gamma_{\alpha\beta})\Big\} + \frac{1}{2}\Big(\hat{\nabla}_\mu + \hat{\Gamma}_{c\mu}{}^{c}\Big)j^\mu + \frac{1}{2}\Big(\hat{\nabla}_a + \hat{\Gamma}_{\alpha a}{}^{\alpha}\Big)j^a, \qquad \text{(B32)}
\end{aligned}$$

respectively, and the scalar curvature R is given by their sum,

$$\begin{aligned}
R &= \gamma^{\mu\nu}\tilde{R}_{\mu\nu} + \phi^{ac}R_{ac} + \frac{1}{4}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b} + \frac{1}{4}\gamma^{\mu\nu}\phi^{ab}\phi^{cd}\Big\{(D_\mu\phi_{ac})(D_\nu\phi_{bd}) \\
&\quad -(D_\mu\phi_{ab})(D_\nu\phi_{cd})\Big\} + \frac{1}{4}\phi^{ab}\gamma^{\mu\nu}\gamma^{\alpha\beta}\Big\{(\partial_a\gamma_{\mu\alpha})(\partial_b\gamma_{\nu\beta}) - (\partial_a\gamma_{\mu\nu})(\partial_b\gamma_{\alpha\beta})\Big\} \\
&\quad +(\hat{\nabla}_\mu + \hat{\Gamma}_{c\mu}{}^{c})j^\mu + (\hat{\nabla}_a + \hat{\Gamma}_{\alpha a}{}^{\alpha})j^a. \qquad \text{(B33)}
\end{aligned}$$

We summarize the notations below:
1. $\tilde{R}_{\mu\nu}$ is defined as

$$\tilde{R}_{\mu\nu} = \hat{\partial}_\mu\hat{\Gamma}_{\alpha\nu}{}^{\alpha} - \hat{\partial}_\alpha\hat{\Gamma}_{\mu\nu}{}^{\alpha} + \hat{\Gamma}_{\mu\beta}{}^{\alpha}\hat{\Gamma}_{\alpha\nu}{}^{\beta} - \hat{\Gamma}_{\beta\alpha}{}^{\beta}\hat{\Gamma}_{\mu\nu}{}^{\alpha}. \qquad \text{(B34)}$$

Notice that the connection coefficients $\hat{\Gamma}_{\mu\nu}{}^{\alpha}$ are given by

$$\hat{\Gamma}_{\mu\nu}{}^{\alpha} = \frac{1}{2}\gamma^{\alpha\beta}\Big(\hat{\partial}_\mu\gamma_{\nu\beta} + \hat{\partial}_\nu\gamma_{\mu\beta} - \hat{\partial}_\beta\gamma_{\mu\nu}\Big), \qquad \text{(B35)}$$

and that $\tilde{R}_{\mu\nu}$ is defined using the metric $\gamma_{\mu\nu}$ of the horizontal space and the horizontal vector fields $\hat{\partial}_\mu$ only. Therefore $\tilde{R}_{\mu\nu}$ is the Ricci tensor of the horizontal space.
2. R_{ac} is defined as

$$R_{ac} = \partial_a\hat{\Gamma}_{bc}{}^{b} - \partial_b\hat{\Gamma}_{ac}{}^{b} + \hat{\Gamma}_{ad}{}^{b}\hat{\Gamma}_{bc}{}^{d} - \hat{\Gamma}_{db}{}^{d}\hat{\Gamma}_{ac}{}^{b}. \qquad \text{(B36)}$$

The connection coefficients $\hat{\Gamma}_{ab}{}^{c}$ of the fibre space N_2 are given by

$$\hat{\Gamma}_{ab}{}^{c} = \frac{1}{2}\phi^{cd}\Big(\partial_a\phi_{bd} + \partial_b\phi_{ad} - \partial_d\phi_{ab}\Big). \qquad \text{(B37)}$$

Therefore R_{ac} is the Ricci tensor of N_2.
3. For completeness, we quote here the definition of the diffN_2-field strength $F_{\mu\nu}{}^{a}$ and the diffN_2-covariant derivative $D_\mu\phi_{ab}$,

$$\begin{aligned}
F_{\mu\nu}{}^{a} &= \partial_\mu A_\nu{}^{a} - \partial_\nu A_\mu{}^{a} - [A_\mu, A_\nu]^a_{\rm L}, \\
D_\mu\phi_{ab} &= \partial_\mu\phi_{ab} - [A_\mu, \phi]_{{\rm L}ab}. \qquad \text{(B38)}
\end{aligned}$$

As is shown in Appendix A, $F_{\mu\nu}{}^{a}$ and $D_\mu\phi_{ab}$ transform covariantly under the diffN_2 transformations.

4.The derivative operators $\hat{\nabla}_\mu$ and $\hat{\nabla}_a$ are the natural derivative operators compatible with $\gamma_{\alpha\beta}$ and ϕ_{bc}. In the horizontal lift basis, the compatibility conditions can be written as

$$\hat{\nabla}_\mu \gamma_{\alpha\beta} = \hat{\partial}_\mu \gamma_{\alpha\beta} - \hat{\Gamma}_{\mu\alpha}{}^{\delta} \gamma_{\delta\beta} - \hat{\Gamma}_{\mu\beta}{}^{\delta} \gamma_{\alpha\delta} = 0, \tag{B39}$$

$$\hat{\nabla}_a \phi_{bc} = \partial_a \phi_{bc} - \hat{\Gamma}_{ab}{}^{d} \phi_{dc} - \hat{\Gamma}_{ac}{}^{d} \phi_{bd} = 0, \tag{B40}$$

respectively. These properties can be also checked using the formulae (B14) and (B20).
5.We also defined j^μ and j^a as

$$j^\mu = \gamma^{\mu\nu} \phi^{ab} D_\nu \phi_{ab}, \tag{B41}$$

$$j^a = \phi^{ab} \gamma^{\mu\nu} \partial_b \gamma_{\mu\nu}. \tag{B42}$$

Thus $\hat{\nabla}_\mu j^\mu$ and $\hat{\nabla}_a j^a$ are given by

$$\hat{\nabla}_\mu j^\mu = \hat{\partial}_\mu j^\mu + \hat{\Gamma}_{\mu\nu}{}^{\mu} j^\nu, \tag{B43}$$

$$\hat{\nabla}_a j^a = \partial_a j^a + \hat{\Gamma}_{ab}{}^{a} j^b, \tag{B44}$$

respectively.

The last two terms containing j^μ and j^a in (B33) turn out to be surface terms in the action integral, as can be seen in what follows. Let us consider the term that contains j^μ. It becomes

$$\begin{aligned} \sqrt{-\gamma}\sqrt{\phi}\,\Big(\hat{\nabla}_\mu + \hat{\Gamma}_{c\mu}{}^{c}\Big) j^\mu &= \sqrt{-\gamma}\sqrt{\phi}\,\Big(\hat{\partial}_\mu j^\mu + \hat{\Gamma}_{\nu\mu}{}^{\nu} j^\mu + \hat{\Gamma}_{c\mu}{}^{c} j^\mu\Big) \\ &= \sqrt{-\gamma}\sqrt{\phi}\,\Big(\partial_\mu j^\mu - A_\mu{}^{a} \partial_a j^\mu + \hat{\Gamma}_{\nu\mu}{}^{\nu} j^\mu + \hat{\Gamma}_{c\mu}{}^{c} j^\mu\Big). \end{aligned} \tag{B45}$$

The first term in the right hand side of (B45) can be written as

$$\sqrt{-\gamma}\sqrt{\phi}\,\partial_\mu j^\mu = -\frac{1}{2}\sqrt{-\gamma}\sqrt{\phi}\,\Big(\gamma^{\alpha\beta}\partial_\mu \gamma_{\alpha\beta} + \phi^{ab}\partial_\mu \phi_{ab}\Big) j^\mu + \partial_\mu\Big(\sqrt{-\gamma}\sqrt{\phi}\, j^\mu\Big), \tag{B46}$$

and the second term becomes

$$\begin{aligned} -\sqrt{-\gamma}\sqrt{\phi}\, A_\mu{}^{a} \partial_a j^\mu = {}& \frac{1}{2}\sqrt{-\gamma}\sqrt{\phi}\, A_\mu{}^{a}\Big(\gamma^{\alpha\beta}\partial_a \gamma_{\alpha\beta} + \phi^{bc}\partial_a \phi_{bc}\Big) j^\mu + \sqrt{-\gamma}\sqrt{\phi}\,(\partial_a A_\mu{}^{a}) j^\mu \\ & -\partial_a\Big(\sqrt{-\gamma}\sqrt{\phi}\, A_\mu{}^{a} j^\mu\Big). \end{aligned} \tag{B47}$$

Using the equations (B24) and (B27), the last two terms in (B45) can be written as

$$\begin{aligned} & \sqrt{-\gamma}\sqrt{\phi}\,\Big(\hat{\Gamma}_{\nu\mu}{}^{\nu} j^\mu + \hat{\Gamma}_{c\mu}{}^{c} j^\mu\Big) \\ ={}& \sqrt{-\gamma}\sqrt{\phi}\,\Big(\frac{1}{2}\gamma^{\alpha\beta}\hat{\partial}_\mu \gamma_{\alpha\beta} + \frac{1}{2}\phi^{ab} D_\mu \phi_{ab}\Big) j^\mu \\ ={}& \sqrt{-\gamma}\sqrt{\phi}\,\Big(\frac{1}{2}\gamma^{\alpha\beta}\hat{\partial}_\mu \gamma_{\alpha\beta} + \frac{1}{2}\phi^{ab}\hat{\partial}_\mu \phi_{ab} - \partial_c A_\mu{}^{c}\Big) j^\mu \\ ={}& \sqrt{-\gamma}\sqrt{\phi}\,\Big(\frac{1}{2}\gamma^{\alpha\beta}\partial_\mu \gamma_{\alpha\beta} - \frac{1}{2}A_\mu{}^{c}\gamma^{\alpha\beta}\partial_c \gamma_{\alpha\beta} + \frac{1}{2}\phi^{ab}\partial_\mu \phi_{ab} - \frac{1}{2}A_\mu{}^{c}\phi^{ab}\partial_c \phi_{ab} - \partial_c A_\mu{}^{c}\Big) j^\mu. \end{aligned} \tag{B48}$$

From these, we find that (B45) is indeed a total divergence,

$$\sqrt{-\gamma}\sqrt{\phi}\,\Big(\hat{\nabla}_\mu+\hat{\Gamma}_{c\mu}{}^{c}\Big)j^\mu=\partial_\mu\Big(\sqrt{-\gamma}\sqrt{\phi}\,j^\mu\Big)-\partial_a\Big(\sqrt{-\gamma}\sqrt{\phi}\,A_\mu{}^{a}j^\mu\Big). \tag{B49}$$

Similarly, we can show that the following is also a total divergence,

$$\sqrt{-\gamma}\sqrt{\phi}\,(\hat{\nabla}_a+\hat{\Gamma}_{\alpha a}{}^{\alpha})j^a=\partial_a\Big(\sqrt{-\gamma}\sqrt{\phi}\,j^a\Big). \tag{B50}$$

Thus the Einstein-Hilbert action integral in this (2,2)-KK formalism can be written as

$$\begin{aligned}
S=\int\!d^2xd^2y\,\sqrt{-\gamma}\sqrt{\phi}\,\Big[&\gamma^{\mu\nu}\tilde{R}_{\mu\nu}+\phi^{ac}R_{ac}+\frac{1}{4}\gamma^{\mu\nu}\gamma^{\alpha\beta}\phi_{ab}F_{\mu\alpha}{}^{a}F_{\nu\beta}{}^{b}\\
&+\frac{1}{4}\gamma^{\mu\nu}\phi^{ab}\phi^{cd}\Big\{(D_\mu\phi_{ac})(D_\nu\phi_{bd})-(D_\mu\phi_{ab})(D_\nu\phi_{cd})\Big\}\\
&+\frac{1}{4}\phi^{ab}\gamma^{\mu\nu}\gamma^{\alpha\beta}\Big\{(\partial_a\gamma_{\mu\alpha})(\partial_b\gamma_{\nu\beta})-(\partial_a\gamma_{\mu\nu})(\partial_b\gamma_{\alpha\beta})\Big\}\Big]\\
&+\int\!d^2xd^2y\,\Big[\partial_\mu\Big(\sqrt{-\gamma}\sqrt{\phi}\,j^\mu\Big)-\partial_a\Big(\sqrt{-\gamma}\sqrt{\phi}\,A_\mu{}^{a}j^\mu\Big)+\partial_a\Big(\sqrt{-\gamma}\sqrt{\phi}\,j^a\Big)\Big].
\end{aligned} \tag{B51}$$

APPENDIX C: THE POSITIVE DEFINITENESS OF κ^2

In this section we will show that κ^2, defined in (4.11) in section IV, is positive definite. To this end, it is convenient to work with the superspace indices $A'=(ac)$, $B'=(bd)$ such that

$$\rho_{A'}=\rho_{ac},\qquad \rho_{B'}=\rho_{bd}. \tag{C1}$$

Define the metric $G_{A'B'}$ on the superspace and its inverse metric $G^{A'B'}$ by

$$G_{A'B'}=\frac{1}{2}(\rho_{ab}\rho_{cd}+\rho_{ad}\rho_{cb}), \tag{C2}$$

$$G^{A'B'}=\frac{1}{2}(\rho^{ab}\rho^{cd}+\rho^{ad}\rho^{cb}), \tag{C3}$$

respectively, so that the following relation

$$G^{A'E'}G_{E'B'}=\delta^{A'}_{B'} \tag{C4}$$

holds. Here the superspace Kronecker delta is defined as

$$\delta^{A'}_{B'}=\frac{1}{2}(\delta^{a}{}_{b}\delta^{c}{}_{d}+\delta^{a}{}_{d}\delta^{c}{}_{b}). \tag{C5}$$

This supermetric raises and lowers the superindices

$$\begin{aligned}
G^{A'B'}\rho_{B'}&=\rho^{A'},\\
G_{A'B'}\rho^{B'}&=\rho_{A'}.
\end{aligned} \tag{C6}$$

Then κ^2 can be written as

$$\begin{aligned}\kappa^2 &= \frac{1}{8}\rho^{ab}\rho^{cd}(D_-\rho_{ac})(D_-\rho_{bd}) \\ &= \frac{1}{8}G^{A'B'}(D_-\rho_{A'})(D_-\rho_{B'}).\end{aligned} \tag{C7}$$

The positive definiteness of κ^2 comes from the positive definiteness of the supermetric $G^{A'B'}$. Let us notice that, when $\rho^{ab} = \delta^{ab}$, this supermetric becomes

$$G^{A'B'} = \mathrm{diag}(+1, +1/2, +1), \tag{C8}$$

which is positive definite. By continuity, it follows that

$$\kappa^2 \geq 0. \tag{C9}$$

REFERENCES

[1] Q.H. Park, Phys. Lett. B **238**, 287 (1990); Int. J. Mod. Phys. A **7**, 1415 (1992).

[2] Y.M. Cho, Q.H. Park, K.S. Soh, and J.H. Yoon, Phys. Lett. B **286**, 251 (1992).

[3] J.H. Yoon, Phys. Lett. B **308**, 240 (1993).

[4] J.H. Yoon,*(1+1)-Dimensional Methods for General Relativity*, in *Directions in General Relativity*, in Proceedings of the 1993 International Symposium, Vol. 2, ed. B.L. Hu and T.A. Jacobson (Cambridge University Press, 1993).

[5] J.H. Yoon, *in preparation.*

[6] H. Bondi, M.G.J. van der Burg, and A.W.K. Metzner, Proc. R. Soc. Lond. A **269**, 21 (1962).

[7] R.K. Sachs, Proc. R. Soc. Lond. A **270**, 103 (1962); J. Math. Phys. **3**, 908 (1962); *Gravitational Radiation*, in *Relativity, Groups and Topology*, The 1963 Les Houches Lectures, ed. B. deWitt and C. deWitt (Gordon and Beach, New York, 1964).

[8] A. Ashtekar, *Asymptotic Quantization*, *Monographs and Textbooks in Physical Sciences* (Bibliopolis, 1986).

[9] A. Ashtekar, J. Bičak, and B. Schmidt, *Asymptotic Structure of Symmetry Reduced General Relativity*, gr-qc/9608042.

[10] A.E. Dominguez, C.N. Kozameh, and M. Ludvigsen, *Superselection Sectors in Asymptotic Quantization of Gravity*, gr-qc/9609071.

[11] H. Friedrich and J.M. Stewart, Proc. R. Soc. Lond. A **385**, 345 (1983).

[12] J.N. Goldberg, Found. Phys. **14**, 1211 (1984); *ibid.* **15**, 439 (1985); *ibid.* **15**, 1075 (1985).

[13] C.G. Torre, Class. Quantum. Grav. **3**, 773 (1986).

[14] Y.M. Cho, J. Math. Phys. **16**, 2029 (1975).

[15] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (Freeman, 1972).

[16] R. Geroch, in *Asymptotic Structure of Space-time*, ed. P. Esposito and L. Witten (Plenum, 1977).

[17] P.R. Garabedian, *Partial Differential Equations* (John Wiley and Sons, 1966).

[18] E.T. Newman and T.W.J. Unti, J. Math. Phys. **3**, 891 (1962).

[19] R.A. d'Inverno and J. Stachel, J. Math. Phys. **19**, 2447 (1978).

[20] R.A. d'Inverno and J. Smallwood, Phys. Rev. D **22**, 1233 (1980).

[21] J. Smallwood, J. Math. Phys. **24**, 599 (1983).

[22] E.T. Newman and K.P Tod, *Asymptotically Flat Space-Times*, in *General Relativity and Gravitation*, Vol.2, ed. A. Held (Plenum Press, 1980).

[23] D. Brill, Ann. Phys. **7**, 466 (1959); Nuovo Cimento Supp. **2**, 1 (1964).

[24] J.H. Yoon, C.M. Kim, and S.K. Oh, *A New Way of Looking at Exact Solutions: Schwartzschild Solution in the (2,2) Kaluza-Klein Formalism of General Relativity*, *in preparation.*

[25] D. Kramer *et al*, *Exact Solutions of Einstein's Field Equations*, ed. E. Schmutzer (Cambridge University Press, 1980).